Asian Journal of
Science, Technology, Engineering, and Art

e-ISSN: 3025-4507
p-ISSN: 3025-5287
Index: Harvard, Boston, Sydney University, Dimensions, Lens, Scilit, Semantic, Google, etc

https://doi.org/10.58578/AJSTEA.v3i2.5206

Gromov - Hausdorff Dimension and Measure of Separable Metric Spaces

Monsuru A. Morawo[1], Hina A. D.[2], Azeez K. Y.[3]
[1]Bells University of Technology, Ogun State, Nigeria; [2]Gombe State University, Gombe, Nigeria; [3]Obafemi Awolowo University, Osun State, Nigeria
morawomonsuruajibola@gmail.com; dhaliyu@gsu.edu.ng

Article Info:

Submitted:	Revised:	Accepted:	Published:
Feb 20, 2025	Mar 6, 2025	Mar 18, 2025	Mar 23, 2025

Abstract

In this paper, we proved that dimension of Gromov – Hausdorff hyperspace of Riemannian manifold depends on the cardinality of measurable Riemannian manifold.

Keywords: Riemannian Manifold, Separable Space, Topological Dimension, Lebesgue Measure, Hausdorff Measure

Introduction

In this section, we are going to discuss about Gromov - Hausdorff dimension of a separable metric space X. The Gromov – Hausdorff dimension of a metric space need not be an integer (Facundo, and Zhengchao, 2021). Though, it can be used to measure the structure of limit spaces of a sequence of Riemannian manifold. The Gromov – Hausdorff

AJSTEA Journal is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License

measure is also called Gromov – Lebesgue measure, in fact, if the subset $Y \subset X$ is a Lebesgue measurable space in Gromov – sense, then it is called Gromov – Lebesgue measurable space. So, for differentiable manifolds, the Gromov – Hausdorff measure is considered as Gromov – Lebesgue measure (Marc, 2009). The Gromov – Hausdorff distance of metric space X depends on the nature of metric space or Riemannian manifold X. If the metric space X is induced with zero dimensional Gromov – Hausdorff measure, Gromov – Hausdorff measure of $X = 0, iff X = \emptyset$, Gromov – Hausdorff measure of $X = n,$ if and only if X is finite and Gromov – Hausdorff measure of $X > 0$ if and only if X is infinite set, (Fremlin, 2000).

Definition of some important terms

Definition 2.1 (Dong and Gabjin, 2000): Let X be a compact metric space. For $\epsilon > 0,$ define $Cov(X, \epsilon)$ as the minimal number of closed ϵ –balls needed to cover X and $Cap(X, \epsilon)$ as the maximal number of disjoint $\epsilon -$ balls in X . $Cov(X, \epsilon)$ is called $\epsilon - covering$ and $Cap(X, \epsilon)$ is called $\epsilon - capacity$ of X .

Definition2.2 (Gromov, 1954)**:** A metric space X is totally bounded if for any $\epsilon > 0,$ there exists a finite number N_ϵ of $\epsilon - balls \{B_i = B(x_i, \epsilon)\}_{i=1}^{N_\epsilon}$ which covers X.

Definition 2.3 (Morawo, *et.al.,* 2024): A topological space X is finite if $\dim(X) < \infty$.

Definition 2.4 (Morawo, *et.al.,* 2024): A metric space X is complete if there is a sequence x_n in X such that $x_n \to L \in X$.

Definition 2.5 (Monsuru, Ahmadu and Balla, 2020): A function $f: X \to Y$ is continuous, if for $O \in Y, f^{-1}(O) \in X$.

Definition 2.6 (Monsuru and Ahmadu, 2020): A topological space X is everywhere dense in Y if and only if $cl(X) = Y$.

Definition 2.7 (Morawo, *et.al.,* 2024): A topological space X is called a Riemannia manifold, if Riemannian metric is defined on it.

Definition 2.8 (Morawo, *et.al.,* 2024): A topological space of infinite dimensional is called an infinite dimensional topological space.

Definition 2.9 (Willard, 2004): A topological space X is separable if it has a countable everywhere dense subset.

Definition 2.10 (Morawo, *et.al.,* 2024): A function $f: X \to Y$ is called $\varepsilon - isometry$ if $|d_Y(f(x) - f(y)) - d_X(x, y)| < \varepsilon$ for every $x, y \in X$.

Definition 2.11 (Morawo , *et.al.*, 2023): A function $f: X \to Y$ is called $\delta - isometry$ if $|d_Y(f(x) - f(y)) - d_X(x, y)| < \delta$ for every $x, y \in X$.

Definition 2.12 (Shanti and Raisinganian, 1965): A function $f: X \to Y$ is called an $isometry$ if $d_Y(f(x) - f(y)) \leq d_X(x, y)$, for every $x, y \in X$.

Definition 2.13 (Morawo, *et.al.,* 2024): A map $i: X \to Y$ is an isometric, if X and Y are isometry space.

Definition 2.14 (Paige, 2022): A topological space X is Hausdorff if for $A, B \in X, A \cap B = \emptyset$.

Methodology

By (Gromov, 1954) and (Olga, G. M., and Peter, W.M. 1991), let X be a separable metric space and p an arbitrary real number, $0 \leq p < \infty$. Given $\epsilon > 0$, let

$\mathcal{GH}_{p.\epsilon}(X) = \frac{\pi^{p/2}}{\Gamma(\frac{p}{2}+1)} inf\{\sum_{i=1}^{\infty} r_i^p : X = \bigcup_{i=1}^{\infty} B(x_i, r_i), r_i \leq \epsilon\}$, where $B(x_i, r_i)$ represents the open ball in X with radius r_i centered at $x_i \in X$ and Γ is called a Gamma function defined by

$\Gamma(t) = \int_0^{\infty} e^{-s} s^{t-1} ds$, (Zorich, 2004)

Note that if $p \geq 2$ is a positive integer, then the constant

$\frac{\pi^{p/2}}{\Gamma(\frac{p}{2}+1)} = \frac{1}{p} vol(S^{p-1}) = \frac{\omega_{p-1}}{p}$, which is exactly equal to $\frac{1}{p} - multiple$ of the volume of round $(p-1) - sphere$. Recall that the volume of a ball of radius r in the Euclidean space $\mathbb{R}^p$ is given by $\frac{1}{p} vol(S^{p-1})$.

Definition 3.1 (Dong and Gabjin, 2000)**:** Let X be a separable metric space and p an arbitrary real number, $0 \leq p < \infty$. The $p-$ dimensional Gromov - Hausdorff measure of X is defined as $$\mathcal{GH}_p(X) = \lim_{\epsilon \to 0} \mathcal{GH}_{p.\epsilon}(X).$$

The Gromov - Hausdorff measure looks like the Lebesgue measure. In fact, if A is a Lebesgue measurable subset of $\mathbb{R}^n$, the $n-$ dimensional Gromov - Hausdorff measure $\mathcal{GH}_n(A)$ is equal to Lebesgue measure of A (Masaki, 2022).

For smooth manifolds, The Gromov - Hausdorff measure can be consider as generalization of the Lebesgue measure (Facundo, 2012). The basic properties are the followings;

Proposition 3.1 (Dong, and Gabjin, 2000)**:** For the zero dimensional $GH-$ measure, we have the following properties;

a. $\mathcal{GH}_0(X) = 0$, if $X = \emptyset$
b. $\mathcal{GH}_0(X) = n$, if X is finite set of n points
c. $\mathcal{GH}_0(X) = \infty$, if X is an infinite set

Proposition 3.2: (Dong, and Gabjin, 2000) If $p < q$, then $\mathcal{GH}_p(X) \geq \mathcal{GH}_q(X)$; in fact, $p < q$ and $\mathcal{GH}_p(X) < \infty$ imply that $\mathcal{GH}_q(X) = 0$.

4.0 Results and Discussion

Now, we are going to make use of **Proposition 3.1** above to provide proof for the **Theorem 4.1** below. Also, note that a metric space is separable if it contains a countable everywhere dense subset, and since every finite countable space is compact, then we have the result below.

Theorem 4.1: Suppose X is a compact separable metric space. Then $\mathcal{GH}_p(X) = 0$ if and only if for each $\epsilon > 0$, there is a finite decomposition of X as:

$X = O_1 \cup O_2 \cup \ldots \cup O_n, O_i = O(x_i, r_i)$ such that

$$\sum_{i=1}^{n} k_i^p < \epsilon.$$

Proof: Suppose $\mathcal{GH}_p(X) = 0$ and given that there is $\epsilon > 0$. By definition separability of a metric space, there is a countable number of balls $O(x_1, k_i^1), O(x_2, k_2^1), \ldots$ such that

$$X =$$

$\bigcup_{i=1}^{\infty} O(x_1, k_i^1)$ *and* $\sum_{i=1}^{\infty} {k^1}_i^p < \frac{\epsilon}{2}$.

It is easy to expand each ball $O(x_i, k_i^1)$ faintly to an open ball $O(x_i, k_i)$ such that

$$k_i^p < k^{1}{}_i^p + \frac{\epsilon}{2^{i+1}}.$$

Provided that metric X is compact, then there exists finite covers $O_1, O_2, \ldots, O_n$, $B_k = O(x_n, k_n)$ of X. Therefore,

$$\sum_{i=1}^{n} k_i^p < \epsilon. \blacksquare$$

In the next result, we are going to make use of **definition 4.2, Lemma 4.2** below and **Proposition 3.2** above to establish the dimension of Gromov – Hausdorff hyperspace and its convergence in the Gromov sense.

Definition 4.2 (Paige, 2022)**:** The Gromov – Hausdorff dimension of an arbitrary separable metric space X is given by

$$dim_{\mathcal{GH}}(X) = inf\{p \geq 0 : \mathcal{GH}_p(X) = 0\} = sup\{p \geq 0 : GH_p(X) > 0\}$$

Note that even if $dim_{\mathcal{GH}}(X) = p$, we have $\mathcal{GH}_p(X) = 0, \infty$ or positive real number. In particular, for a smooth Riemannian $n-$ manifold (M^g, g), we have $dim_{\mathcal{GH}}(X) = n$ and $\mathcal{GH}_n(M) = vol(M, g)$.

Lemma 4.2 (Dong and Gabjin, 2000)**:** For a compact metric space X and $\epsilon > 0$, $Cov(X, 2\epsilon) \leq Cap(X, \epsilon)$.

Theorem 4.2: Given positive real number $k, P > 0$ and a natural number $n \in \mathbb{N}$, let (M_i, g_i) be a sequence of Riemannian n – manifold satisfying $Ric(M_i) \geq -(n-1)k$, $diam(M_i) \leq P$. Suppose X_l is the Gromov – Hausdorff hyperspace limit of the sequence (M_i, g_i). Then $dim_{\mathcal{GH}} X_l \leq n$.

Proof: By **proposition 3.2**, it suffices to show that $\mathcal{GH}_n(X_l) < \infty$. Recall that X_l is a compact inner metric space with $diam(X_l) \leq P$. For every $\epsilon > 0$, recall also that $Cov(X, \epsilon)$ denotes the smallest number of closed $\epsilon - balls\ in\ X$ which cover X.

Also, provided that $Cov(X, \epsilon)$ is continuous in the Gromov Hausdorff sense by **Lemma 4.2**, we have

$Cov(X_l, \epsilon) \leq Cov\left(M_i, \epsilon - \frac{3}{\delta_i}\right), \delta_i \to 0 \; as \; i \to \infty$, for i sufficiently large. The volume comparison theorem together with $diam(M_i) \leq P$, implies that $Cov(M_i, \epsilon)$ depends only on $\epsilon, k \; and \; P$. Thus, we have

$$\mathcal{GH}_n(X_l) = \lim_{\epsilon \to 0} \frac{\omega_{n-1}}{n} \epsilon^n Cov(X_l, \epsilon) \leq \lim_{\epsilon \to 0} \left\{ \lim_{i \to \infty} sup \frac{\omega_{n-1}}{n} \epsilon^n cov\left(M_i, \epsilon - \frac{3}{\delta_i}\right) \right\}$$

$$= \lim_{i \to \infty} sup \left(\frac{\omega_{n-1}}{n} \epsilon^n cov\left(M_i, \epsilon - \frac{3}{\delta_i}\right) \right)$$

$$= \lim_{i \to \infty} sup \, \mathcal{GH}_n(M_i) \leq V_k(P) < \infty.$$

By volume comparison theorem, the last inequality follows. Hence,

$GH_n(M_i) = vol(M_i)$, for a compact smooth Riemannian manifold■

In the next result, we are going to make use of **Proposition 4.1 and Lemma 4.2** below to establish that in a Gromov sense, dimension of hyperspace associated with Riemannian n – manifold is finite dimensional.

Proposition 4.1 (Gromov, 1954)**:** For positive real numbers k and D and a natural number $n \in \mathbb{N}$, let (M_i, g_i) be a sequence of Riemannian n – manifold satisfying $Ric(M_i) \geq -(n-1)k, \; diam(M_i) \leq D$. Suppose X_l is the Gromov – Hausdorff hyperspace limit of the sequence (M_i, g_i). Then every point $x \in X_l$ has a tangent cone $\mathcal{GH} \lim_{r_i}(X, x, r_i) \equiv T_x X_l$

Lemma 4.2 (Facundo, 2021)**:** For any fixed $\epsilon > 0 \; and \; R > \epsilon$, the function $f: X \to N(\epsilon, R, X)$ is almost continuous in the Gromov – Hausdorff distance.

Theorem 4.3: Given positive real numbers $k, v, D > 0$ and a natural number $n \in \mathbb{N}$, let (M_i, g_i) be a sequence of Riemannian n – manifold satisfying

$Ric(M_i) \geq -(n-1)k \, , vol(M_i) \geq v \; and \; diam(M_i) \leq D$. Suppose X_l is the Gromov – Hausdorff hyperspace limit of the sequence (M_i, g_i). Then $dim_{GH} X_l = n$ and the tangent cone $T_x X_l$ at any point $x \in X_l$ has Gromov – Hausdorff dimension n, i.e

$dim_{\mathcal{GH}} T_x X_l = dim_{\mathcal{GH}} \lim_{r \to \infty}(X_l, x, r.d) = n$, for any point $x \in X_l$.

Proof: Since $vol(M_i) \geq v$, one has $dim_{GH}(X_l) \geq n$ and it follows from the **Theorem 4.2** that $dim_{\mathcal{GH}}(X_l) = n$. It remains to prove that $dim_{\mathcal{GH}} T_x X_l = n$. It is known from the

volume condition that for each point $x \in X_l, T_xX$ is a metric cone , i.e , metrically cone on tangent space and so it becomes a length space. As in the proof of **Proposition 3.1,** one has $N(\epsilon, R(X_l, rd)) \leq C\left(\frac{R}{\epsilon}\right)^n$ and so $dim_{\mathcal{GH}} T_x X_l \leq n.$ To show the inequality, it is enough to verify that for any $\alpha > 0, \mathcal{GH}_{n-\alpha}(T_x X_l \cap B(R=1) = \infty,$ where we designate $B(R=1)$ as the unit ball in the tangent cone $T_x X_l$ centred at x. Then, we claim that:

$N(\epsilon, R(X_l, rd)) \leq C^1\left(\frac{R}{\epsilon}\right)^n$, for some positive constant $C^1 = C^1(k, v, D, n) = \infty.$

Note that, for $N = N(\epsilon, R, X_l)$,

$volB(R) \leq \sum_{i=1}^{N} volB(R = 2\epsilon) \leq N.\max\{volB(R = 2\epsilon)\}.$ ----------(*)

Provided that N is almost continuous by **Lemma 4.2** above, we have

$N\left(\epsilon, R(X_l, rd)\right) = N\left(\frac{\epsilon}{r}, \frac{R}{r}, (X, d)\right) \geq C_1.N(\frac{\epsilon}{r}, \frac{R}{r}, M_i) \geq \frac{volB\left(\frac{R}{r}\right)}{\max\left\{volB\left(\frac{2\epsilon}{r}\right)\right\}}$, from equation (*) above.

Note that,

$volB\left(\frac{2\epsilon}{r}\right) \leq V_k\left(\frac{2\epsilon}{r}\right) \leq C\left(\frac{R}{r}\right)^n$, for $r >> 1$ adequately large. So,

$\frac{volB\left(\frac{2\epsilon}{r}\right)}{\max\left\{volB\left(\frac{2\epsilon}{r}\right)\right\}} \geq C_2 \frac{volB\left(\frac{2\epsilon}{r}\right)}{\left(\frac{2\epsilon}{r}\right)^n} \geq C_2 \frac{vol(M)}{V_k(D)} \frac{V_k\left(\frac{R}{r}\right)}{\left(\frac{2\epsilon}{r}\right)^n}$, from relative volume comparison theorem

$\geq C_2 \frac{v}{V_k(D)} \frac{\left(\frac{R}{r}\right)^n}{\left(\frac{2\epsilon}{r}\right)^n} = C_1\left(\frac{2\epsilon}{r}\right)^n,$ where $C_1 = C_1(k, v, D, n).$

Finally, choose $\sigma - net$ in $B(R=1)$ so that balls of radius σ are disjoint and $2\sigma - balls$ cover $B(R=1)$.

Then for any $\alpha > 0$, we have

$\sum \frac{\omega_{n-1}}{n}\sigma^{n-\alpha} = \frac{\omega_{n-1}}{n}\sigma^{n-\alpha}N(\sigma, 1, T_x X_l) = \frac{\omega_{n-1}}{n}\sigma^{n-\alpha},$

Hence, $dim_{\mathcal{GH}} T_x X_l \geq n$ by definition, this completes the proof■

Conclusion

In this work, we proved that Dimension of Gromov – Hausdorff hyperspace of Riemannian manifold depends on the cardinality of Riemannian manifold.

References

Dong, C., and Gabjin, Y. (2000). "Gromov – Hausdorff Topology and its application to Riemannia Manifold.

Fremlin, D.H. (2019). Measure theory. *Volume 5. University of Essex*.

Facundo, M and Zhengchao (2021). Charaterization of Gromov – type geodesics. *Department of Mathematics and Computer Science and Engineering, The Ohio State University.*

Facundo, M. (2012). Some properties of Gromov – Hausdorff distance. *Discrete compute Geom. 48. 416 – 440.*

Fell, J. (1962). A Hausdorff topology for the closed subset of a locally compact and non – Hausdorff spaces. *Proc. Amer.Math.Soc.13, 472 – 476*.

Gromov, M. (1954). Metric structures for Riemannian and Non – Riemannian Spaces.

Lee, G.T. (2018). Abstract algebra: An introductory course. *Springer Undergraduate Mathematics Series, Canada. Doi: 10.1007/978 – 3 – 319*.

Masaki, M. (2022) "Mean Hausdorff Dimension of some infinite dimensional fractals"2020 Mathematics subject classification.

Monsuru, A.M., Ahmadu, K., and Balla, M. Y. (2020). On the comparative study of compactness and some of its relative notions in Metric and Topological spaces. *International Journal of scientific and Research Publication, Vol. 10 Issue 10. 659 – 665*.

Murray. (2021). The 54th Spring Topology and Dynamic Conference. *Murray State University. USA.*

Marc, C. (2009). Volume and Topology.

Morawo, M.A, Danyaro, M.L, Baily, A.S and Azeez, K.Y. (2024) Gromov – Hausdorff Convergence and Topological stability for Actions on Wasserstein Hyperspaces. Journal of Applied Science and Environmental Management. 28(11B). 3815 – 3822.

Morawo, M.A, Azeez, K.Y, Danyaro, M.L, Baily, A. S and Mohammed, A.B. (2024) Comparative Study of Some Topological Properties on Hyperspace of Convex Bodies and Wasserstein hyperspace Associated with Riemannian Manifold. Journal of Applied Science and Environmental Management. 28(12). 4051 – 4056.

Monsuru A Morawo, Mshelia, I. B, Manjak, N.H and Hina, A.D. (2023) Topological properties of some hyperspaces of convex bodies Associated with Riemannian Manifold. International Journal of Advanced in Engineering and Management. 5(3). Pp: 1494 – 1499.

Shanty, N and Raisinganian, M.D. (1965) Element of Real analysis. S. Chand & Company LTD. Ram Nager, New Delhi – 110 055.

Olaf, M. (2021) "Gheeger – Gromov compactness theorem for Manifold with boundary. Adv. Math. 224 – 240.

Olga, G.M., and Peter W.M. (1991). The Riemannian manifold of all Riemannia metrics. *Quarterly Oxford Journal of Mathematics 42(183 - 202).*

Paige, D. (2022). Hausdorff Dimension and Projection

Willard, S. (2004). General topology. *Additson – Wesley.*

Zorich, V.A (2004) Mathematical Analysis II. Springer.