

RATIONAL CONTRACTION IN METRIC SPACE AND COMMON FIXED POINT THEOREMS

Surendra Kumar Tiwari & Jayant Prakash Ganvir

Dr. C. V. Raman University, Bilaspur, India

sk10tiwari@gmail.com; jayantganveer@gmail.com

Article Info:

Submitted:	Revised:	Accepted:	Published:
Jan 1, 2024	Jan 12, 2024	Jan 15, 2024	Jan 18, 2024

Abstract

The study of contraction mappings in fixed-point theory is a fascinating and crucial field of mathematics. The concept of contraction plays a vital role in proving the existence and uniqueness of fixed points. Banach's contraction theory offers a fixed point theorem that is widely accepted as unique in most analyses. By using rational expressions in metric spaces, we can achieve unique results in general contraction mapping. These results are based on several innovative ideas stemming from the latest research. The delivered results upgrade and federate many existing outcomes on the topic in the literature Bhardwaj, R. et al. (2007) Chouhan et al. (2014) and Garg and Priyanka (2016). Also gives some suitable examples for verifying our results.

Keywords: Complete Metric space, Fixed point, Common Fixed Point, Rational expressions

INTRODUCTION

The literature of fixed point (FP) for contraction mapping embarked upon Banach contraction principle (BCP) in 1922 and known as the Banach fixed point Theorem (BFP). BCP is stated as follows:

“A single valued contractive type mapping on a complete metric space has a unique FP.”

The fixed point theorem can be used to prove that a differential equation, an integral equation, or a partial differential equation has a unique solution. They are utilized in the investigation of problems related to the optimal control of these systems.

Outcomes of FP (fixed point) treatment with general contractive conditions relating to rational type expression are also interesting. Some notorious results from this track are included (see [Das and Gupta (1975), Khan, M.S.(1976), Jaggi, D.S.(1977), Fisher, B. (1978), Jaggi and Das (1980), Bhardwaj et al. (2007), Dutta and Chaudhary (2008), Jleli, et al.(2012), Shrivastava, et al.(2013), Karapinar,E., Shatanawi, W. and Tas, K (2013), Saluja et al.(2015), Garg, M.(2015), Garg, M. and Priyanka(2016), Karapinar,E., et al. (2016), Fulga, A.(2018), Gupta, S. and Garg, A.K., (2019), Fulga, A. (2020)]). It has been extended and generalized various types of contraction in different metric spaces.

Dewangan,S. and Tiwari, S.K.(2020), extended, generalized and improved the result of Shrivastava et al (2014) and established a unique fixed point theorem for expansive mapping in metric space satisfying rational expression . In 2021, Rao S. and Kalyani, obtained fixed point results partially ordered metric space with rational expression. Subsequently, Rao, N.S. et al. (2021), demonstrated result of fixed point of monotone functions in ordered metric space.

Furthermore, Raji, M. and Adegboye, F. (2022), proved the existence and uniqueness of some fixed point for nonlinear contractive mappings with rational expressions in the context of metric space endowed with partial order. Most recently, Chaodary et al. (2023) established a result in the fixed point theory for multivalued Mizoguchi-Takahashi type rational contraction in metric space.

In this manuscript, we extend, generalize, and improve earlier work by Bhardwaj, R. et al. (2007), Chouhan et al. (2014), and Garg and Priyanka (2016) to determine in common fixed-point theorems for rational contraction on metric space.

PRELIMINARIES NOTES

We recall the definition of metric space and their properties as follows:

Definition 1 [Shrivastava et al. (2014.)]: Infer that $\mathfrak{S} \neq \phi$ and $d: \mathfrak{S} \times \mathfrak{S} \rightarrow [0, \infty)$ be a mapping that known as metric on $\mathfrak{S}$, if satisfies the following conditions:

- (i) $d(u, v) \geq 0 \forall u, v \in \mathfrak{S}$;
- (ii) $d(u, v) = 0 \Leftrightarrow u = v \forall u, v \in \mathfrak{S}$;
- (iii) $d(u, v) = d(v, u) \forall u, v \in \mathfrak{S}$;
- (iv) $d(u, v) \leq d(u, w) + d(w, v)$, for all $u, v, w \in \mathfrak{S}$.

Then, we can refer to the pair $(\mathfrak{S}, d)$ as a metric space. .

Definition 2[Shrivastava et al. (2014.)]: Assume that $\{u_n\}$ is a sequence referred to Cauchy sequence in $(\mathfrak{S}, d)$, even if $\varepsilon > 0 \exists n_0 \in \mathbb{N}$, like that, $\forall m, n > n_0$

$$d(u_n, u_m) < \varepsilon. i. e. \lim_{n \rightarrow \infty} d(u_n, u_m) < \varepsilon.$$

Definition 3 [Shrivastava et al. (2014.)]: The sequence $\{u_n\}$ is convergent to $u \in \mathfrak{S}$ in $(\mathfrak{S}, d)$, if $\lim_{n \rightarrow \infty} d(u_n, u) = 0$, or $u_n \rightarrow u$.

Definition 4[Shrivastava et al. (2014.)]: when everything the Cauchy sequences converges in $(\mathfrak{S}, d)$ then metric space $(\mathfrak{S}, d)$ is so called complete.

MAIN RESULTS

Theorem 5. Consider that, $(\mathfrak{S}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{S}, \partial)$, where $(\mathfrak{S}, \partial)$ is a complete metric space. Take over, $j, k: \mathfrak{S} \rightarrow \mathfrak{S}$ be a continuous self mapping such that $v, \xi \in \mathfrak{S}$, $v \neq \xi$ there exist $\theta_1, \theta_2, \theta_3, \theta_4, \theta_5 \in [0, 1)$ with $\theta_1 + \theta_2 + \theta_3 + \theta_4 + 2\theta_5 < 2$, satisfies the following condition

$$\begin{aligned} \partial(jv, k\xi) \leq & \theta_1 \frac{\partial(v, jv) \cdot \partial(\xi, k\xi) + \partial(v, k\xi) \cdot \partial(\xi, jv)}{\partial(v, jv) + \partial(\xi, k\xi) + \partial(v, k\xi) + \partial(\xi, jv)} + \theta_2 \frac{\partial(v, jv) \cdot \partial(\xi, jv) + \partial(\xi, k\xi) \cdot \partial(v, k\xi)}{\partial(v, jv) + \partial(\xi, jv) + \partial(\xi, k\xi) + \partial(v, k\xi)} \\ & + \theta_3 \frac{\partial(v, jv) \cdot \partial(v, jv) + \partial(v, k\xi) \cdot \partial(\xi, jv)}{\partial(v, jv) + \partial(v, jv) + \partial(v, k\xi) + \partial(\xi, jv)} + \theta_4 \frac{\partial(v, jv) \cdot \partial(v, k\xi) + \partial(\xi, jv) \cdot \partial(\xi, k\xi)}{\partial(v, jv) + \partial(v, k\xi) + \partial(\xi, jv) + \partial(\xi, k\xi)} \\ & + \theta_5 \partial(v, \xi). \end{aligned} \quad (5.1)$$

For $v_0 \in \mathcal{H}$, suppose $\{v_k\}_{k=0}^\infty \subset \mathfrak{J}$ and defined by $v_{2k+1} = jv_{2k}$ and $v_{2k+2} = \ell v_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to j and ℓ . Then there exist $v_k \rightarrow v^*$, $\lim_{k \rightarrow \infty} j^k v = v^* = \lim_{k \rightarrow \infty} \xi$. Then v^* is a unique common fixed point.

Proof: Let $v_0 \in \mathcal{H}$, we define the iterative sequence $\{v_{2k}\}$ in $\mathfrak{J}$ defined as

$$v_{2k+1} = jv_k = j^{2k+1}v_0,$$

And

$$v_{2k+2} = \ell v_{2k+1} = \ell^{2k+2}v_0, \text{ for every } k \geq 1.$$

Now put $v = v_{2k}$ and $\xi = v_{2k-1}$ in (5.1) we have

$$\begin{aligned} \partial(v_{2k+1}, v_{2k}) &= \partial(jv_{2k}, \ell v_{2k-1}) \\ &\leq \theta_1 \frac{\partial(v_{2k}, jv_{2k}).\partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_{2k}, \ell v_{2k-1}).\partial(v_{2k-1}, jv_k)}{\partial(v_{2k}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_{2k}, \ell v_{2k-1}) + \partial(v_{2k-1}, jv_k)} \\ &\quad + \theta_2 \frac{\partial(v_{2k}, jv_{2k}).\partial(v_{2k-1}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1}).\partial(v_{2k}, \ell v_{2k-1})}{\partial(v_{2k}, jv_{2k}) + \partial(v_{2k-1}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_k, \ell v_{2k-1})} \\ &\quad + \theta_3 \frac{\partial(v_{2k}, jv_{2k}).\partial(v_{2k}, jv_k) + \partial(v_{2k}, \ell v_{2k-1}).\partial(v_{2k-1}, jv_{2k})}{\partial(v_{2k}, jv_{2k}) + \partial(v_{2k}, jv_{2k}) + \partial(v_k, \ell v_{2k-1}) + \partial(v_{k-1}, jv_k)} \\ &\quad + \theta_4 \frac{\partial(v_{2k}, jv_{2k}).\partial(v_k, \ell v_{2k-1}) + \partial(v_{2k-1}, jv_{2k}).\partial(v_{2k-1}, \ell v_{2k-1})}{\partial(v_{2k}, jv_{2k}) + \partial(v_{2k}, \ell v_{2k-1}) + \partial(v_{2k-1}, jv_{2k}) + \partial(v_{k-1}, \ell v_{2k-1})} \\ &\quad + \theta_5 \partial(v_{2k}, v_{2k-1}). \\ &= \theta_1 \frac{\partial(v_{2k}, v_{2k+1}).\partial(v_{2k-1}, v_{2k}) + \partial(v_{2k}, v_{2k}).\partial(v_{2k-1}, v_{2k+1})}{\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k}) + \partial(v_{2k}, v_{2k}) + \partial(v_{2k-1}, v_{2k+1})} \\ &\quad + \theta_2 \frac{\partial(v_{2k}, v_{2k+1}).\partial(v_{2k-1}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k}).\partial(v_{2k}, v_{2k})}{\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k}) + \partial(v_{2k}, v_{2k})} \\ &\quad + \theta_3 \frac{\partial(v_{2k}, v_{2k+1}).\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k}, v_{2k}).\partial(v_{2k-1}, v_{2k+1})}{\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k}, v_{2k+1}) + \partial(v_{2k}, v_{2k}) + \partial(v_{2k-1}, v_{2k+1})} \\ &\quad + \theta_4 \frac{\partial(v_{2k}, v_{2k+1}).\partial(v_{2k}, v_{2k}) + \partial(v_{2k-1}, v_{2k+1}).\partial(v_{2k-1}, v_{2k})}{\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k}, v_{2k}) + \partial(v_{2k-1}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k})} \\ &\quad + \theta_5 \partial(v, v_{k-1}). \\ &\leq \left(\frac{\theta_1}{2} + \frac{\theta_2}{2} + \frac{\theta_3}{2}\right) \partial(v_{2k}, v_{2k+1}) + \left(\frac{\theta_4}{2} + \theta_5\right) \partial(v_{2k-1}, v_{2k}). \end{aligned}$$

Therefore,

$$\partial(v_{2k+1}, v_{2k}) \leq \pi \partial(v_{2k-1}, v_{2k}).$$

Where $\pi = \frac{\left(\frac{\theta_4}{2} + \theta_5\right)}{\left(1 - \frac{\theta_1}{2} - \frac{\theta_2}{2} - \frac{\theta_3}{2}\right)} < 1$, but $\theta_1 + \theta_2 + \theta_3 + \theta_4 + 2\theta_5 < 2$.

So, continuing this process, we get

$$\partial(v_{2k+1}, v_{2k}) \leq \pi^{2k} \partial(v_0, v_1).$$

Now for $l > k$ and using the triangle inequality, we obtain

$$\begin{aligned} \partial(v_{2k}, v_{2l}) &\leq \partial(v_{2k}, v_{2k+1}) + \partial(v_{2k+1}, v_{2k+2}) + \dots + \partial(v_{2l-1}, v_{2l}) \\ &\leq \pi^{2k} \partial(v_0, v_1) + \pi^{2k+1} \partial(v_0, v_1) + \dots + \pi^{2k+2l-1} \partial(v_0, v_1) \\ &\leq (\pi^{2k} + \pi^{2k+1} + \dots + \pi^{2k+2l-1}) \partial(v_0, v_1) \\ &\leq \frac{\pi^{2k}}{1-\pi} \partial(v_0, v_1) \rightarrow \partial(v_k, v_l) \rightarrow 0, k, l \rightarrow \infty. \end{aligned}$$

Hence in $\mathfrak{S}$, $\{v_{2k}\}$ is a

Cauchy sequence. Due to the fact that $(\mathfrak{S}, \partial)$ is a complete metric spaces, $\exists$ a point $v^* \in \mathfrak{S}$

Such that $v_{2k} \rightarrow v^*$ as $k \rightarrow \infty$. Again, since j is a continuous such that $j(v) =$

$j(\lim_{k \rightarrow \infty} v_{2k}) = \lim_{k \rightarrow \infty} jv_{2k} = \lim_{k \rightarrow \infty} v_{2k+1} = v^*$. Hence v^* is fixed point of j in $\mathfrak{S}$. Similarly, we can show that $k(\xi) = v^*$. Since $j(v) = v^* = k(\xi)$. Therefore, v^* is a common fixed point of j and k .

Now we will show that v^* is a unique common fixed point of j and k in $\mathfrak{S}$. Suppose ξ^* is another common fixed point of j and k in $\mathfrak{S}$. then we have

$$\begin{aligned} \partial(v^*, \xi^*) &= \partial(jv^*, k\xi^*) \\ &\leq \theta_1 \frac{\partial(v^*, jv^*) \cdot \partial(\xi^*, k\xi^*) + \partial(v^*, k\xi^*) \cdot \partial(\xi^*, jv^*)}{\partial(v^*, jv^*) + \partial(\xi^*, k\xi^*) + \partial(v^*, k\xi^*) + \partial(\xi^*, jv^*)} \\ &\quad + \theta_2 \frac{\partial(v^*, jv^*) \cdot \partial(\xi^*, jv^*) + \partial(\xi^*, k\xi^*) \cdot \partial(v^*, k\xi^*)}{\partial(v^*, jv^*) + \partial(\xi^*, jv^*) + \partial(\xi^*, k\xi^*) + \partial(v^*, k\xi^*)} \\ &\quad + \theta_3 \frac{\partial(v^*, jv^*) \cdot \partial(v^*, jv^*) + \partial(v^*, k\xi^*) \cdot \partial(\xi^*, jv^*)}{\partial(v^*, jv^*) + \partial(v^*, jv^*) + \partial(v^*, k\xi^*) + \partial(\xi^*, jv^*)} \\ &\quad + \theta_4 \frac{\partial(v^*, jv^*) \cdot \partial(v^*, k\xi^*) + \partial(\xi^*, jv^*) \cdot \partial(\xi^*, k\xi^*)}{\partial(v^*, jv^*) \cdot \partial(v^*, k\xi^*) + \partial(\xi^*, jv^*) \cdot \partial(\xi^*, k\xi^*)} \\ &\quad + \theta_5 \partial(v^*, \xi^*). \\ &= \theta_1 \frac{\partial(v^*, v^*) \cdot \partial(\xi^*, \xi^*) + \partial(v^*, \xi^*) \cdot \partial(\xi^*, v^*)}{\partial(v^*, v^*) + \partial(\xi^*, \xi^*) + \partial(v^*, \xi^*) + \partial(\xi^*, v^*)} \\ &\quad + \theta_2 \frac{\partial(v^*, v^*) \cdot \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*) \cdot \partial(v^*, \xi^*)}{\partial(v^*, v^*) + \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*) + \partial(v^*, \xi^*)} \\ &\quad + \theta_3 \frac{\partial(v^*, v^*) \cdot \partial(v^*, v^*) + \partial(v^*, \xi^*) \cdot \partial(\xi^*, v^*)}{\partial(v^*, v^*) + \partial(v^*, v^*) + \partial(v^*, \xi^*) + \partial(\xi^*, v^*)} \\ &\quad + \theta_4 \frac{\partial(v^*, v^*) \cdot \partial(v^*, \xi^*) + \partial(\xi^*, v^*) \cdot \partial(\xi^*, \xi^*)}{\partial(v^*, v^*) + \partial(v^*, \xi^*) + \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*)} \\ &\quad + \theta_5 \partial(v^*, \xi^*). \end{aligned}$$

Thus $\partial(v^*, \xi^*) \leq \left[\frac{\theta_1}{2} + \frac{\theta_3}{2} + \frac{\theta_4}{2} + \theta_5 \right] \partial(v^*, \xi^*) \rightarrow 0$, which is a contradiction, because $\theta_1 + \theta_2 + \theta_3 + \theta_4 + 2\theta_5 < 2$. So, $\partial(v^*, \xi^*) = 0$. Implies that $v^* = \xi^*$, thus, v^*

is a unique common fixed point of j and k in $\mathfrak{J}$. The theorem's proof approach has become done.

Example 6: Let $\mathfrak{J} = [0, 1]$ with partial order " $\leq$ " and let $j, k: \mathfrak{J} \rightarrow \mathfrak{J}$ be defined by

$$j(v) = \begin{cases} \frac{1}{2} & \text{if } v \in [0, \frac{1}{4}) \\ v - \frac{1}{4} & \text{if } v \in [\frac{1}{4}, 1) \end{cases} = k(\xi)$$

To show that j and k satisfies the contractive condition 51) of theorem (5), for $v = \frac{1}{2}$,

$\xi = \frac{1}{6}$, $\theta_1 = \theta_2 = \theta_3 = \theta_4 = 1$. Then we have

$$jv = \frac{1}{4}, k\xi = \frac{1}{2}, \partial(v, jv) = \frac{1}{4}, \partial(\xi, k\xi) = \frac{1}{3}, \partial(v, k\xi) = 0, \partial(\xi, jv) = \frac{1}{12}, \\ \partial(v, \xi) = \frac{1}{3} \text{ and } \partial(jv, k\xi) = \frac{1}{4}.$$

Thus, from (5.1), we get

$$\begin{aligned} \frac{1}{4} &= \partial(jv, k\xi) \\ &\leq \frac{2}{3} + \frac{1}{32} + \frac{3}{28} + \frac{1}{24} + \frac{1}{3} \theta_5. \\ &= \frac{448+21+72+28}{672} + \frac{1}{3} \theta_5 \\ \frac{1}{4} &= \frac{569}{672} + \frac{1}{3} \theta_5. \end{aligned}$$

It follows that, $\theta_5 \geq \frac{401}{672}$. Hence all the conditions of Theorem 3.1 are satisfied and $\frac{1}{2}$ is a unique common fixed point of j and k .

Remark: If $\theta_1 = \theta_2 = \theta_3 = \theta_4 = 0$ and $\theta_5 = \alpha$, then the theorem reduce to Banach, S. (1922).

Theorem 7: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $j, k: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $v, \xi \in \mathfrak{J}$, $v \neq \xi$ there exist $\theta_1, \theta_2, \theta_3, \theta_4 \in [0, 1)$ with $2\theta_1 + 2\theta_2 + \theta_4 < 2$, satisfies the following condition

$$\begin{aligned} \partial(jv, k\xi) &\leq \theta_1 \partial(v, \xi) + \theta_2 \frac{\partial(v, jv) \cdot \partial(\xi, k\xi) + \partial(v, k\xi) \cdot \partial(\xi, jv)}{\partial(v, \xi)} \\ &+ \theta_3 \frac{\partial(v, jv) [\partial(v, jv) + \partial(\xi, k\xi)]}{\partial(v, \xi) + \partial(\xi, jv) + \partial(\xi, k\xi)} \end{aligned}$$

$$+ \theta_4 \frac{\partial(v, jv) \cdot \partial(\xi, jv) + \partial(\xi, \ell\xi) \cdot \partial(v, \ell\xi)}{\partial(v, jv) + \partial(\xi, jv) + \partial(\xi, \ell\xi) + \partial(v, \ell\xi)}. \quad (7.1)$$

For $v_0 \in \mathcal{H}$, suppose $\{v_k\}_{k=0}^\infty \subset \mathcal{H}$ and defined by $v_{2k+1} = jv_{2k}$ and $v_{2k+2} = \ell v_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to j and ℓ . Then there exist $v_k \rightarrow v^*$, $\lim_{k \rightarrow \infty} j^k v = v^* = \lim_{k \rightarrow \infty} \xi$. Then v^* is a unique common fixed point.

Proof: : Let $v_0 \in \mathfrak{J}$, we define the iterative sequence $\{v_{2k}\}$ in $\mathfrak{J}$ defined as

$$v_{2k+1} = jv_k = j^{2k+1}v_0,$$

And

$$v_{2k+2} = \ell v_{2k+1} = \ell^{2k+2}v_0, \text{ for every } k \geq 1.$$

Now put $v = v_{2k}$ and $\xi = v_{2k-1}$ in (7.1) we have

$$\begin{aligned} \partial(v_{2k+1}, v_{2k}) &= \partial(jv_k, \ell v_{2k-1}) \\ &\leq \theta_1 \partial(v_{2k}, v_{2k-1}) \\ &+ \theta_2 \frac{\partial(v_{2k}, jv_{2k}) \cdot \partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_{2k}, \ell v_{2k-1}) \cdot \partial(v_{2k-1}, jv_{2k})}{\partial(v_{2k}, v_{2k-1})} \\ &+ \theta_3 \frac{\partial(v_{2k}, \ell v_{2k-1}) [\partial(v_{2k}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1})]}{\partial(v_{2k}, v_{2k-1}) + \partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_{2k-1}, jv_{2k})} \\ &+ \theta_4 \frac{\partial(v_{2k}, jv_k) \cdot \partial(v_{2k-1}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1}) \cdot \partial(v_{2k}, \ell v_{2k-1})}{\partial(v_{2k}, jv_{2k}) + \partial(v_{2k-1}, jv_{2k}) + \partial(v_{2k-1}, \ell v_{2k-1}) + \partial(v_{2k}, \ell v_{2k-1})} \\ &= \theta_1 \partial(v_{2k}, v_{2k-1}) \\ &+ \theta_2 \frac{\partial(v_{2k}, v_{2k+1}) \cdot \partial(v_{2k-1}, v_{2k}) + \partial(v_{2k}, v_{2k}) \cdot \partial(v_{2k-1}, v_{2k+1})}{\partial(v_{2k}, v_{2k-1})} \\ &+ \theta_3 \frac{\partial(v_{2k}, v_{2k}) [\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k})]}{\partial(v_{2k}, v_{2k-1}) + \partial(v_{2k-1}, v_{2k}) + \partial(v_{2k-1}, v_{2k+1})} \\ &+ \theta_4 \frac{\partial(v_{2k}, v_{2k+1}) \cdot \partial(v_{2k-1}, v_{2k+1}) + \partial(v_{2k}, v_{2k}) \cdot \partial(v_{2k-1}, v_{2k+1})}{\partial(v_{2k}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k+1}) + \partial(v_{2k-1}, v_{2k}) + \partial(v_{2k}, v_{2k})} \\ &\leq \theta_1 \partial(v_{2k}, v_{2k-1}) + (\theta_2 + \frac{\theta_4}{2}) \partial(v_{2k}, v_{2k-1}). \end{aligned}$$

Therefore,

$$\partial(v_{2k+1}, v_{2k}) \leq \Psi \partial(v_{2k}, v_{2k-1}).$$

Where $\Psi = \frac{\theta_1}{(1 - \theta_2 - \frac{\theta_4}{2})} < 1$, but $2\theta_1 + 2\theta_2 + \theta_4 < 2$.

So, continuing this process, we get

$$\partial(v_{2k+1}, v_{2k}) \leq \Psi^k \partial(v_0, v_1).$$

Now for $l > k$ and using the triangle inequality, we obtain

$$\partial(v_{2k}, v_{2l}) \leq \partial(v_{2k}, v_{2k+1}) + \partial(v_{2k+1}, v_{2k+2}) + \dots + \partial(v_{2l-1}, v_{2l})$$

$$\begin{aligned} &\leq \Psi^{2k} \partial(v_0, v_1) + \Psi^{2k+1} \partial(v_0, v_1) + \dots + \Psi^{2k+2l-1} \partial(v_0, v_1) \\ &\leq (\Psi^{2k} + \Psi^{2k+1} + \dots + \Psi^{2k+2l-1}) \partial(v_0, v_1) \\ &\leq \frac{\Psi^{2k}}{1-\Psi} \partial(v_0, v_1) \rightarrow 0. \end{aligned}$$

Hence $\partial(v_{2k}, v_{2l}) \rightarrow 0$, $k, l \rightarrow \infty$. Hence in $\mathfrak{J}$, $\{v_{2k}\}$ is a Cauchy sequence. Due to the fact that $(\mathfrak{J}, \partial)$ is a complete cone metric spaces, $\exists v^* \in \mathfrak{J}$ Such that $v_k \rightarrow v^*$ as $k \rightarrow \infty$. Again, since j and k are continuous such that

$$\begin{aligned} j(v) &= j(\lim_{k \rightarrow \infty} v_{2k}) = \lim_{k \rightarrow \infty} jv_{2k} = \lim_{k \rightarrow \infty} v_{2k+1} = v^* \text{ and} \\ k(v) &= jk(\lim_{k \rightarrow \infty} v_{2k}) = \lim_{k \rightarrow \infty} kv_{2k} = \lim_{k \rightarrow \infty} v_{2k+2} = v^*. \end{aligned}$$

Hence v^* is common fixed point of j and k in $\mathfrak{J}$.

Now we will show that v^* is a unique common fixed point of j and k in $\mathfrak{J}$.

Suppose ξ^* is another common fixed point of j and k in $\mathfrak{J}$. Then from (7.1) we have

$$\begin{aligned} \partial(v^*, \xi^*) &= \partial(jv^*, k\xi^*) \\ &\leq \theta_1 \partial(v^*, \xi^*) + \theta_2 \frac{\partial(v^*, jv^*) \cdot \partial(\xi^*, k\xi^*) + \partial(v^*, k\xi^*) \cdot \partial(\xi^*, jv^*)}{\partial(v^*, \xi^*)} \\ &\quad + \theta_3 \frac{\partial(v^*, jv^*) [\partial(v^*, jv^*) + \partial(\xi^*, k\xi^*)]}{\partial(v^*, \xi^*) + \partial(\xi^*, jv^*) + \partial(\xi^*, k\xi^*)} + \theta_4 \frac{\partial(v^*, jv^*) \cdot \partial(\xi^*, jv^*) + \partial(\xi^*, k\xi^*) \cdot \partial(v^*, k\xi^*)}{\partial(v^*, jv^*) + \partial(\xi^*, jv^*) + \partial(\xi^*, k\xi^*) + \partial(v^*, k\xi^*)} \\ &\leq \theta_1 \partial(v^*, \xi^*) + \theta_2 \frac{\partial(v^*, v^*) \cdot \partial(\xi^*, \xi^*) + \partial(v^*, \xi^*) \cdot \partial(\xi^*, v^*)}{\partial(v^*, \xi^*)} \\ &\quad + \theta_3 \frac{\partial(v^*, v^*) [\partial(v^*, v^*) + \partial(\xi^*, \xi^*)]}{\partial(v^*, \xi^*) + \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*)} + \theta_4 \frac{\partial(v^*, v^*) \cdot \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*) \cdot \partial(v^*, \xi^*)}{\partial(v^*, v^*) + \partial(\xi^*, v^*) + \partial(\xi^*, \xi^*) + \partial(v^*, \xi^*)}. \end{aligned}$$

Thus

$\partial(v^*, \xi^*) \leq (\theta_1 + \theta_2) \partial(v^*, \xi^*) \rightarrow 0$, which is a contradiction, because $2\theta_1 + 2\theta_2 + \theta_4 < 2$, So, $\partial(v^*, \xi^*) = 0$. Implies that $v^* = \xi^*$, thus, v^* is a unique common fixed point of j and k in $\mathfrak{J}$.

Example 8: Let $\mathfrak{J} = \{\frac{1}{4}, \frac{1}{2}, 1\}$ with partial order " $\leq$ " and $\partial(v, \xi) = |v - \xi|$ be a cone metric space on $\mathfrak{J}$. Assume that $j, k : \mathfrak{J} \rightarrow \mathfrak{J}$ be defined by

$$j(v) = \begin{cases} v^2, & \text{if } v \in [0, \frac{1}{2}] \\ 1 - v, & \text{if } v \in (\frac{1}{2}, 1] \\ \frac{v}{2}, & \text{if } v \in (1, \infty) \end{cases} = k(\xi)$$

To show that j and k are satisfies the contractive condition (7.1) of theorem (7), for $v = \frac{1}{2}, \xi = 1, \theta_2 = \theta_3 = \theta_4 = 1$. Then we have

$$\begin{aligned} jv &= \frac{1}{4}, k\xi = \frac{1}{2}, \partial(v, jv) = \partial\left(\frac{1}{2}, j\frac{1}{2}\right) = \frac{1}{4}, \\ \partial(\xi, k\xi) &= \partial(1, k1) = \frac{1}{2}, \\ \partial(v, k\xi) &= \partial\left(\frac{1}{2}, k1\right) = 0, \\ \partial(\xi, jv) &= \partial\left(1, j\frac{1}{2}\right) = \frac{3}{4}, \\ \partial(v, \xi) &= \partial\left(\frac{1}{2}, 1\right) = \frac{1}{2} \text{ and } \partial(jv, k\xi) = \partial\left(j\frac{1}{2}, k1\right) = \frac{1}{4}. \end{aligned}$$

Thus, from (3.1.1), we get

$$\begin{aligned} \frac{1}{4} &= \partial(jv, k\xi) \\ &\leq \frac{1}{2}\theta_1 + \frac{1}{4} + 0 + \frac{1}{8} \\ &= \frac{1}{2}\theta_1 + \frac{3}{8}. \end{aligned}$$

It follows that, $\theta_1 \leq \frac{1}{4}$. Hence all the conditions of Theorem 7 are satisfied and j and k has a unique fixed point $v = \frac{1}{4} = \xi$.

Corollary 9: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $j, k: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $v, \xi \in \mathfrak{J}$, $v \neq \xi$ there exist $\theta_1, \theta_2 \in [0, 1)$ with $\theta_1 + \theta_2 < 2$, satisfies the following condition

$$\partial(jv, k\xi) \leq \theta_1 \partial(v, \xi) + \theta_2 \frac{\partial(v, jv) \cdot \partial(\xi, k\xi) + \partial(v, k\xi) \cdot \partial(\xi, jv)}{\partial(v, \xi)} \quad (9.1)$$

For $v_0 \in \mathfrak{J}$, suppose $\{v_k\}_{k=0}^\infty \subset \mathfrak{J}$ and defined by $v_{2k+1} = jv_{2k}$ and $v_{2k+2} = kv_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to j and k . Then there exist $v_k \rightarrow v^*, \lim_{k \rightarrow \infty} j^k v = v^* = \lim_{k \rightarrow \infty} \xi$. Then v^* is a unique common fixed point.

Theorem 10: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $\mathcal{L}_1, \mathcal{L}_2: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $q, r \in \mathfrak{J}$, $q \neq r$ there exist $i_1, i_2 \in [0, 1)$ with $2i_1 + i_2 < 1$, satisfies the following condition

$$\partial(\mathcal{L}_1 q, \mathcal{L}_2 r) \leq i_1 \max \left\{ \frac{\frac{\partial(q, \mathcal{L}_1 q) \cdot \partial(r, \mathcal{L}_2 r) + \partial(q, \mathcal{L}_2 r) \cdot \partial(r, \mathcal{L}_1 q)}{\partial(q, r)}, \frac{\partial(q, \mathcal{L}_1 q) \cdot \partial(q, \mathcal{L}_2 r) + \partial(r, \mathcal{L}_2 r) \cdot \partial(r, \mathcal{L}_1 q)}{\partial(q, r)} \right\} \\ + i_2 \partial(q, r)$$

For $q_0 \in \mathfrak{J}$, suppose $\{q_k\}_{k=0}^\infty \subset \mathfrak{J}$ and defined by $q_{2k+1} = \mathcal{L}_1 q_{2k}$ and $q_{2k+2} = \mathcal{L}_2 q_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to $\mathcal{L}_1$ and $\mathcal{L}_2$. Then there exist $q_k \rightarrow q^*$, $\lim_{k \rightarrow \infty} \mathcal{L}_1^k q = q^* = \lim_{k \rightarrow \infty} \mathcal{L}_2 r$. Then q^* is a unique common fixed point.

Proof: Let $q_0 \in \mathfrak{J}$, we define the iterative sequence $\{q_{2k}\}$ in $\mathfrak{J}$ defined as

$$q_{2k+1} = \mathcal{L}_1 q_{2k} = \mathcal{L}_1^{2k+1} q_0,$$

and

$$q_{2k+2} = \mathcal{L}_2 q_{2k+1} = \mathcal{L}_2^{2k+2} q_0, \text{ for every } k \geq 1.$$

Now put $q = q_{2k}$ and $r = q_{2k-1}$ in (9.1) we have

$$\begin{aligned} \partial(q_{2k+1}, q_{2k}) &= \partial(\mathcal{L}_1 q_{2k}, \mathcal{L}_2 q_{2k-1}) \\ &\leq i_1 \max \left\{ \frac{\frac{\partial(q_{2k}, \mathcal{L}_1 q_{2k}) \cdot \partial(q_{2k-1}, \mathcal{L}_2 q_{2k-1}) + \partial(q_{2k}, \mathcal{L}_2 q_{2k-1}) \cdot \partial(q_{2k-1}, \mathcal{L}_1 q_{2k})}{\partial(q_{2k}, q_{2k-1})}, \frac{\partial(q_{2k}, \mathcal{L}_1 q_{2k}) \cdot \partial(q_{2k}, \mathcal{L}_2 q_{2k-1}) + \partial(q_{2k-1}, \mathcal{L}_2 q_{2k-1}) \cdot \partial(q_{2k}, \mathcal{L}_1 q_{2k})}{\partial(q_{2k}, q_{2k-1})} \right\} \\ &\quad + i_2 \partial(q_{2k}, q_{2k-1}). \\ &\leq i_1 \max \left\{ \frac{\frac{\partial(q_{2k}, q_{2k+1}) \cdot \partial(q_{2k-1}, q_{2k}) + \partial(q_{2k}, q_{2k}) \cdot \partial(q_{2k-1}, q_{2k+1})}{\partial(q_{2k}, q_{2k-1})}, \frac{\partial(q_{2k}, q_{2k+1}) \cdot \partial(q_{2k}, q_{2k}) + \partial(q_{2k-1}, q_{2k}) \cdot \partial(q_{2k}, q_{2k+1})}{\partial(q_{2k}, q_{2k-1})} \right\} \\ &\quad + i_2 \partial(q_{2k}, q_{2k-1}). \\ &\leq i_1 \max\{\partial(q_{2k}, q_{2k+1}), \partial(q_{2k}, q_{2k+1})\} + i_2 \partial(q_{2k}, q_{2k-1}). \\ &= i_1 \partial(q_{2k}, q_{2k+1}) + i_2 \partial(q_{2k}, q_{2k-1}). \end{aligned}$$

Thus

$$\partial(q_{2k+1}, q_{2k}) \leq \frac{i_2}{1-i_1} \partial(q_{2k}, q_{2k-1}). \text{ Since } w = \frac{i_2}{1-i_1} < 1.$$

Therefore, $\partial(q_{2k+1}, q_{2k}) \leq w \partial(q_{2k}, q_{2k-1})$. Continuing this process, we get

$$\partial(q_{2k+1}, q_{2k}) \leq w^k \partial(q_0, q_1).$$

Now for $l > k$ and using the triangle inequality, we obtain

$$\partial(q_{2k}, q_{2l}) \leq \frac{w^{2k}}{1-w} \partial(q_0, q_1) \rightarrow 0.$$

Hence $\partial(q_{2k}, q_{2l}) \rightarrow 0, k, l \rightarrow \infty$. Hence in $\mathfrak{S}$, $\{q_{2k}\}$ is a Cauchy sequence. Due to the fact that $(\mathfrak{S}, \partial)$ is a complete cone metric spaces, $\exists q^* \in \mathfrak{S}$ Such that $q_{2k} \rightarrow q^*$ as $k \rightarrow \infty$. Again, since $\mathcal{L}_1$ and $\mathcal{L}_2$ are continuous such that

$$\mathcal{L}_1(q) = \mathcal{L}_1(\lim_{k \rightarrow \infty} q_{2k}) = \lim_{k \rightarrow \infty} \mathcal{L}_1 q_{2k} = \lim_{k \rightarrow \infty} q_{2k+1} = q^* \text{ and}$$

$$\mathcal{L}_2(q) = \mathcal{L}_2(\lim_{k \rightarrow \infty} q) = \lim_{k \rightarrow \infty} \mathcal{L}_2 q_{2k} = \lim_{k \rightarrow \infty} q_{2k+2} = q^*.$$

Since $\mathcal{L}_1(q) = q^* = \mathcal{L}_2(q)$. Therefore, q^* is common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{S}$.

Now we will show that q^* is a unique common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{S}$.

Suppose r^* is another common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{S}$. Then from (9.1) we have

$$\begin{aligned} \partial(q^*, r^*) &= \partial(\mathcal{L}_1 q^*, \mathcal{L}_2 r^*) \\ &\leq i_1 \max \left\{ \frac{\partial(q^*, \mathcal{L}_1 q^*) \cdot \partial(r^*, \mathcal{L}_2 r^*) + \partial(q^*, \mathcal{L}_2 r^*) \cdot \partial(r^*, \mathcal{L}_1 q^*)}{\partial(q^*, r^*)}, \frac{\partial(q^*, \mathcal{L}_1 q^*) \cdot \partial(q^*, \mathcal{L}_2 r^*) + \partial(r^*, \mathcal{L}_2 r^*) \cdot \partial(r^*, \mathcal{L}_1 q^*)}{\partial(q, r)} \right\} + i_2 \partial(q^*, r^*) \\ &\leq (i_1 + i_2) \partial(q^*, r^*) \end{aligned}$$

$\partial(q^*, r^*) \leq (i_1 + i_2) \partial(q^*, r^*) \rightarrow 0$, which is a contradiction, because $2i_1 + i_2 < 1$.

So, $\partial(q^*, r^*) = 0$. Implies that $q^* = r^*$, thus, q^* is a unique common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{S}$. This completes the proof of the theorem.

Lemma 11[Jovanovic, M. et al. (2010): Let $\{q_k\}$ be a sequence in a metric type space $(\mathfrak{S}, d)$ such that

$$d(q_k, q_{k+1}) \leq \sigma d(q_{k-1}, q_k)$$

for some $0 < \sigma < 1$, and each $k = 1, 2, 3 \dots$, then $\{q_k\}$ is a Cauchy sequence in $(\mathfrak{S}, d)$.

Proposition 12: Presume that $(\mathfrak{S}, d)$ is a metric space and $\mathcal{T}: \mathfrak{S} \rightarrow \mathfrak{S}$ be a given function and $\{q_{2k}\}$ be Picard sequence of initial point $q_0 \in \mathfrak{S}$ satisfying the following condition:

$$d(q_{2k}, q_{2k+1}) \leq \frac{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + n}{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + n} d(q_{2k-1}, q_{2k})$$

.... (12.1) where $m, n \in \mathbb{R}^+$ such that $n < m$. Then $\{q_k\}$ is a Cauchy sequence.

Proof: Suppose that $q_0 \in \mathfrak{S}$ be an arbitrary point and let $\{q_{2k}\}$ be sequence of initial point q_0 . If $q_{2k_0} = q_{2k_0-1}$ for some $q_0 \in \mathbb{N}$, then q_{k_0} is a fixed point of $\mathcal{T}$ and so $\{q_{2k}\}$ is a Cauchy sequence. Let us suppose that (12.1) satisfied for the sequence $\{q_{2k}\}$. If $q_{2k_0} \neq q_{2k_0-1}$ for some $k \in \mathbb{N}$, from (12.1) we deduce that the sequence $\{d(q_{2k}, q_{2k+1})\}$ is decreasing. Thus there exist a non negative real number r such

that $d(q_{2k}, q_{2k+1}) \rightarrow r$. Then, we claim that $r = 0$. If $r > 0$, on taking limit as $k \rightarrow \infty$ on both side of (10.1), we get

$$r \leq \frac{r+\dots+r+n}{r+\dots+r+m} r < r, \text{ which is a contradiction. It follows that } r = 0.$$

Now show that $\{q_{2k}\}$ is a Cauchy sequence. Suppose $\sigma \in [0,1)$. Since $r = 0$, there exist $k(\sigma) \in \mathbb{N}$ such that

$$\begin{aligned} & d(q_{2k}, q_{2k+1}) \\ & \leq \frac{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + n}{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + n} \leq \sigma. \end{aligned}$$

For all $k \geq k(\sigma)$.

This implies that $d(q_{2k}, q_{2k+1}) \leq d(q_{2k-1}, q_{2k})$ for all $k \geq k(\sigma)$ and by Lemma 11 $\{q_{2k}\}$ is a Cauchy sequence. The proof that $\{q_{2k}\}$ is a Cauchy sequence if (10.1) holds is the same.

Theorem 13: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $\mathcal{L}_1, \mathcal{L}_2: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $q, r, \in \mathfrak{J}$, $q \neq r$ there exist $\alpha \in [0,1)$ satisfies the following condition

$$\begin{aligned} d(\mathcal{L}_1 q, \mathcal{L}_2 r) & \leq \frac{d(q, \mathcal{L}_2 r) + d(q, \mathcal{L}_2(\mathcal{L}_2 r)) + d(r, \mathcal{L}_1 q) + d(r, \mathcal{L}_1(\mathcal{L}_1 q)) + n}{d(q, \mathcal{L}_1 q) + d(q, \mathcal{L}_1(\mathcal{L}_1 q)) + d(r, \mathcal{L}_2 r) + d(r, \mathcal{L}_2(\mathcal{L}_2 r)) + m} d(q, r) \\ & \quad + \alpha d(r, \mathcal{L}_1 q) \dots \end{aligned} \quad (13.1)$$

For $q_0 \in \mathfrak{J}$, suppose $\{q_{2k}\}_{k=0}^{\infty} \subset \mathfrak{J}$ and defined by $q_{2k+1} = \mathcal{L}_1 q_{2k}$ and $q_{2k+2} = \mathcal{L}_2 q_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to $\mathcal{L}_1$ and $\mathcal{L}_2$. Then there exist $q_{2k} \rightarrow q^*$, $\lim_{k \rightarrow \infty} \mathcal{L}_1^k q = q^* = \lim_{k \rightarrow \infty} r$. Then q^* is a unique common fixed point.

Proof: Let $q_0 \in \mathfrak{J}$, we define the iterative sequence $\{q_{2k}\}$ in $\mathfrak{J}$ defined as

$$q_{2k+1} = \mathcal{L}_1 q_{2k} = \mathcal{L}_1^{2k+1} q_0,$$

and

$$q_{2k+2} = \mathcal{L}_2 q_{2k+1} = \mathcal{L}_2^{2k+2} q_0, \text{ for every } k \geq 1.$$

Now put $q = q_{2k-1}$ and $r = q_{2k}$ in (11.1) we have

$$\begin{aligned} & d(q_{2k}, q_{2k+1}) = d(\mathcal{L}_1 q_{2k-1}, \mathcal{L}_2 q_{2k}) \\ & \leq \frac{d(q_{2k-1}, \mathcal{L}_2 q_{2k}) + d(q_{2k-1}, \mathcal{L}_2(\mathcal{L}_2 q_{2k})) + d(q_{2k}, \mathcal{L}_1 q_{2k-1}) + d(q_{2k}, \mathcal{L}_1(\mathcal{L}_1 q_{2k-1})) + n}{d(q, \mathcal{L}_1 q_{2k-1}) + d(q, \mathcal{L}_1(\mathcal{L}_1 q_{2k-1})) + d(q_{2k}, \mathcal{L}_2 q_{2k}) + d(q_{2k}, \mathcal{L}_2(\mathcal{L}_2 q_{2k})) + m} d(q_{2k-1}, q_{2k}) \\ & \quad + \alpha d(q_{2k}, \mathcal{L}_1 q_{2k-1}) \end{aligned}$$

$$\begin{aligned}
 &\leq \frac{d(q_{2k-1}, q_{2k+1}) + d(q_{2k-1}, L_2(q_{2k+1})) + d(q_{2k}, q_{2k}) + d(q_{2k}, L_1(q_{2k})) + n}{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, L_1(q_{2k})) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, L_2(q_{2k+1})) + m} d(q_{2k-1}, q_{2k}) \\
 &\quad + \alpha d(q_{2k}, q_{2k}) \\
 &\leq \frac{d(q_{2k-1}, q_{2k+1}) + d(q_{2k-1}, q_{2k+2}) + d(q_{2k}, q_{2k+1}) + n}{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + m} d(q_{2k-1}, q_{2k}) \\
 &\leq \frac{[d(q_{2k-1}, q_{2k}) + d(q_{2k}, q_{2k+2})] + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + n}{d(q_{2k-1}, q_{2k}) + d(q_{2k-1}, q_{2k+1}) + d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + m} d(q_{2k-1}, q_{2k}).
 \end{aligned}$$

That is, condition (13.1) holds for the sequence $\{q_{2k}\}$. Then, by proposition (12.1), $\{q_{2k}\}$ is a Cauchy sequence. Since $\mathfrak{J}$ is a complete metric space, the sequence $\{q_{2k}\}$ converges to some $x \in \mathfrak{J}$. Now, we prove that x is a common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$. Using (13.1) with $q = q_{2k}$ and $r = x$, then we get

$$\begin{aligned}
 d(q_{2k+1}, L_2 x) &= d(L_1 q_{2k}, L_2 x) \\
 &\leq \left[\frac{d(q_{2k}, L_2 x) + d(q_{2k}, L_2(L_2 x)) + d(x, L_1 q_{2k}) + d(x, L_1(L_1 q_{2k})) + n}{d(q_{2k}, L_1 q_{2k}) + d(q_{2k}, L_1(L_1 q_{2k})) + d(x, L_2 x) + d(x, L_2(L_2 x)) + m} \right] d(q_{2k}, x) \\
 &\quad + \alpha d(x, L_1 q_{2k}) \\
 &\leq \left[\frac{d(q_{2k}, L_2 x) + d(q_{2k}, L_2(L_2 x)) + d(x, q_{2k+1}) + d(x, q_{2k+2}) + n}{d(q_{2k}, q_{2k+1}) + d(q_{2k}, q_{2k+2}) + d(x, L_2 x) + d(x, L_2(L_2 x)) + m} \right] d(q_{2k}, x) \\
 &\quad + \alpha d(x, q_{2k+1})
 \end{aligned}$$

On taking limit as $k \rightarrow \infty$ on both sides, we get

$$\begin{aligned}
 d(x, L_2 x) &\leq \left[\frac{d(x, L_2 x) + d(q_{2k}, L_2(L_2 x)) + n}{d(x, L_2 x) + d(x, L_2(L_2 x)) + m} \right] \lim_{n \rightarrow \infty} d(q_{2k}, x) \\
 &\quad + \lim_{n \rightarrow \infty} \alpha d(x, q_{2k+1}) = 0.
 \end{aligned}$$

This implies that $d(x, L_2 x) = 0 \Rightarrow x = L_2 x$. Similarly, we can show that $x = L_1 x$. Since $L_1 x = x = L_2 x$. Therefore, x is common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$.

Now we will show that x is a unique common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{J}$.

Suppose y is another common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{J}$. Then from (13.1) with $q = x$ and $r = y$ we get

$$\begin{aligned}
 d(x, y) &= d(L_1 x, L_2 y) \\
 &\leq \left[\frac{d(x, L_2 y) + d(y, L_1 x) + d(x, L_2(L_2 y)) + d(y, L_1(L_1 x)) + n}{d(x, L_1 x) + d(x, L_1(L_1 x)) + d(y, L_2 y) + d(y, L_2(L_2 y)) + m} \right] d(x, y) + \alpha d(y, L_1 x) \\
 &= \frac{4 d(x, y) + n}{m} + \alpha d(y, x) \rightarrow 0.
 \end{aligned}$$

Therefore, $d(x, y) = 0 \Rightarrow x = y$. Thus x is unique common fixed point of $\mathcal{L}_1$ and $\mathcal{L}_2$ in $\mathfrak{J}$. This completes the proof of the theorem.

If $n = m = 0$ in Theorem 13, we get the following Corollary

Corollary 14: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $\mathcal{L}_1, \mathcal{L}_2: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $q, r, \in \mathfrak{J}$, $q \neq r$ there exist $\alpha \in [0, 1)$ satisfies the following condition

$$d(\mathcal{L}_1 q, \mathcal{L}_2 r) \leq \frac{d(q, \mathcal{L}_2 r) + d(q, \mathcal{L}_2(\mathcal{L}_2 r)) + d(r, \mathcal{L}_1 q) + d(r, \mathcal{L}_1(\mathcal{L}_1 q))}{d(q, \mathcal{L}_1 q) + d(q, \mathcal{L}_1(\mathcal{L}_1 q)) + d(r, \mathcal{L}_2 r) + d(r, \mathcal{L}_2(\mathcal{L}_2 r))} d(q, r) + \alpha d(r, \mathcal{L}_1 q) \dots$$

For $q_0 \in \mathfrak{J}$, suppose $\{q_{2k}\}_{k=0}^{\infty} \subset \mathfrak{J}$ and defined by $q_{2k+1} = \mathcal{L}_1 q_{2k}$ and $q_{2k+2} = \mathcal{L}_2 q_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to $\mathcal{L}_1$ and $\mathcal{L}_2$. Then there exist $q_{2k} \rightarrow q^*$, $\lim_{k \rightarrow \infty} \mathcal{L}_1^k q = q^* = \lim_{k \rightarrow \infty} r$. Then q^* is a unique common fixed point.

If $n = m = 1$ in Theorem 13, we get the following Corollary

Corollary 15: Consider that, $(\mathfrak{J}, \leq)$ be a ordered set of partially and let ∂ is a metric on $(\mathfrak{J}, \partial)$, where $(\mathfrak{J}, \partial)$ is a complete metric space. Take over, $\mathcal{L}_1, \mathcal{L}_2: \mathfrak{J} \rightarrow \mathfrak{J}$ be a continuous self mapping such that $q, r, \in \mathfrak{J}$, $q \neq r$ there exist $\alpha \in [0, 1)$ satisfies the following condition

$$d(\mathcal{L}_1 q, \mathcal{L}_2 r) \leq \frac{d(q, \mathcal{L}_2 r) + d(q, \mathcal{L}_2(\mathcal{L}_2 r)) + d(r, \mathcal{L}_1 q) + d(r, \mathcal{L}_1(\mathcal{L}_1 q)) + 1}{d(q, \mathcal{L}_1 q) + d(q, \mathcal{L}_1(\mathcal{L}_1 q)) + d(r, \mathcal{L}_2 r) + d(r, \mathcal{L}_2(\mathcal{L}_2 r)) + 1} d(q, r) + \alpha d(r, \mathcal{L}_1 q) \dots$$

For $q_0 \in \mathfrak{J}$, suppose $\{q_{2k}\}_{k=0}^{\infty} \subset \mathfrak{J}$ and defined by $q_{2k+1} = \mathcal{L}_1 q_{2k}$ and $q_{2k+2} = \mathcal{L}_2 q_{2k+1}$ $k = 0, 1, 2, \dots$ be the Picard iteration association to $\mathcal{L}_1$ and $\mathcal{L}_2$. Then there exist $q_{2k} \rightarrow q^*$, $\lim_{k \rightarrow \infty} \mathcal{L}_1^k q = q^* = \lim_{k \rightarrow \infty} r$. Then q^* is a unique common fixed point.

CONCLUSION

We have investigated and extended the current literature of Bhardwaj R. et al. (2007), Chouhan et al. (2014), and Garg and Priyanka (2016) and obtained unique common fixed point theorems for rational type contraction in metric spaces. Theorem 5 with Example 6, Theorem 7 with Example 8 and Corollary 9, Theorem 10 with Proposition 12, and Theorem 13 with Corollaries 14 and 15 are our results.

REFERENCES

- Banach, S. Sur les operations dans les ensembles abstraits et leur applications aux equation Integrals, *Fundamental Mathematicae*, 3(7), (1922), 133-181.
<https://doi.org/10.4064/fm-3-1131-181>.
- Chandok, S. , Choudhury, B.S., and Metiya, N. Fixed point results in ordered metric spaces for rational type expressions with auxiliary functions, *Journal of the Egyptian Mathematical Society*, 23 (2015), 95-101.
<http://dx.doi.org/10.1016/j.joems.2014.02.002>.
- Choudhury, B.S. , Metiya, N., Chakraborty, P. and Sunirmal Kundu , Multivalued mizoguchi-takahashi type rational contraction in relational metric spaces, *Surveys in Mathematics and its Applications*, Volume 18 (2023), 59 – 72.
<https://www.utgjiu.ro/math/sma>.
- Dass, B.K. and Gupta, S. An Extension of Banach Contraction Principle through Rational Expression, *Indian J. of Pure and Applied Mathematics*, 6(1975), 1455-1458.
- Shakuntala Dewangan and Tiwari, S.K. Expansive Mapping in Metric Spaces and Fixed Point Theorem, *International Journal of Mathematical Archive-* 11(12),(2020), 1-7.
- Dutta, P. N. and Choudhary, B.S. (2008), A Generalization of Contraction Principle in Metric Spaces, *Fixed Point Theory and Appl.*, Art. Id. 406368, 8 pages.
<https://doi.org/10.1155/2008/406368>
- Khan, M. S. A fixed point theorems for metric spaces, *Rendicanti Dell'I Istitute Mathematica Dell' Universita, Trieste Int. J. Math.* 8 (1976), 69-72.
- Jaggi, D.S. Some unique fixed point theorems, *Indian J. Pure Appl. Math*, vol. 8, pp. (1977) 223-230.
- Fisher, B. A fixed point theorem for compact metric spaces, *Publ. Math. Debrecen* 25 (1978), 193-194. <https://doi.org/10.5486/pmd.1978.25.3-4.01>
- Jaggi, D.S. and. Dass,B.K. An extension of Banach contraction theorem through rational expression, *Bull. Cal. Math.*(1980), 261-266.
- Bhardwaj, R., Rajput,S.S., and Yadav, R.N. Application of fixed point theory in metric spaces, *Thai. J. of Mathematics*, 2(2007) 253-259. [www.math.science, cmu.ac.th/thaijournal](http://www.math.science.cmu.ac.th/thaijournal)
- Karapinar,E., Shatanawi, W. and Tas, K. Fixed point theorem on partial metric spaces involving rational expressions, *Miskolc Mathematical Notes*, 14(2013),no:1, 135-142. DOI:[10.18514/MMN.2013.471](https://doi.org/10.18514/MMN.2013.471)
- . Karapinar,E., Roldan A., and Sadarangani,K. Existence and uniqueness of best proximity points under rational contractivity conditions, *Mathematica Slovaca*, 66,(6) (2016) 1427-1442. <https://doi.org/10.1515/ms-2016-0234>.
- Karapinar, E., Dehici,A., and. Redjel, N. On some fixed points of $\alpha - \psi$ contractive mappings with rational expressions *J. Nonlinear Sci. Appl.*, 10 (2017), 1569-1581. doi:[10.22436/jnsa.010.04.23](https://doi.org/10.22436/jnsa.010.04.23).
- Fulga, A. , Fixed point theorems in rational via Suzuki approach results in nonlinear analysis,1(10 (2018) 19-29.
- Fulga, A. On (ψ, ϕ) - rational contractions, *Symmetry*, (2020), 1-13.
<https://doi.org/10.3390/sym12050723>.
- Garg, M. (2015), A New Type Generalized Fixed Point Theorem in a Complete Metric Space, *Ultra. Sci.*, 27(3), (2015), 1-2.
- Garg, M. and Priyanka A New Generalization of Fixed Point Theorem in a Complete Metric Space, *Mathematica Moravica.*, 28(3), (2016),179-182.

- Gupta, S. and Garg, A.K., Fixed Point Theorems on Complete Metric Space, International Journal on Emerging Technologies 10(2b), (2019),109-112
- Proinov, P.D., Fixed point theorems for generalized contractive mappings in metric spaces, J. fixed point theory and App. (2020) 22: 21, DOI:10.1007/s11784-020-0756-
- Raji, M. and Adegbeye, F. Some Fixed point theorems for rational contractions in partially ordered metric spaces, Research Square, (2022), 1-11. DOI: <https://doi.org/10.21203/rs.3.rs-1818692/v1>
- Rao, N. S. and Kalyani, K. (2021), Generalized fixed point results of rational type Contractions in partially ordered metric spaces, BMC Res Notes 14:390, (2021), 1-13. <https://doi.org/10.1186/s13104-021-05801-7>.
- Rao.N.S. , Kalyani, K. and Gemechu,T. Fixed Point Theorems in Partially Ordered Metric Spaces with Rational Expressions, Information Sciences Letters,10(3),(2021), 451-460. doi:10.18576/isl/100309
- Shrivastava, R., Dubey, K.R. and Tiwari, P. Common Fixed Point Theorems in Complete Metric Space, Advances in applied sciences research, 4(6), (2013), 82-89. www.pelagaearesearchlibrary.com
- Shrivastava, R., Sharma, M. and Bhardwaj, R., Fixed point theorems in complete metric space. Journal of multidisciplinary Engineering Science and Technology, 1(4), (2014), 21-27.