African Journal of Sciences and Traditional Medicine

ISSN : 1595-7942

Index: Harvard, Boston, Sydney University, Dimensions, Lens, ResearchGet Scilit, Semantic, Google Scholar, Base etc

https://doi.org/10.58578/AJSTM.v3i3.11654

Effect of Polyherbal Formulated Teaonsperm Count, Motility and Viability in Wistar Rats

Dickson Onyenmulo Uwaya[1], Christian Chukwuebuka Udemmadu[2], Juilet N. Olayinka[3]

[1,2]University of Benin, Benin City, Nigeria

[3]Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria

dickson.uwaya@uniben.edu; udickson4christ@yahoo.com

Article Info:

Submitted:	Revised:	Accepted:	Published:
Jun 19, 2026	Jul 17, 2026	Jul 29, 2026	Aug 3, 2026

Abstract

Polyherbal formulations have attracted increasing scientific interest because their combined constituents may produce synergistic therapeutic effects. This study evaluated the effects of an aqueous polyherbal tea extract on sperm count, motility, viability, and testicular oxidative stress markers in male Wistar rats. Twenty rats were randomly allocated into five groups of four animals: a distilled-water control group, a carbamazepine-treated group receiving 20 mg/kg, a sildenafil citrate-treated group receiving 5 mg/kg, and two treatment groups receiving 5 or 10 mg/kg of the polyherbal tea extract orally for 21 days. Following treatment, the animals were sacrificed by cervical dislocation, and sperm count, progressive and non-progressive motility, and viability were assessed. Testicular tissue homogenates were also analysed for superoxide dismutase, catalase, glutathione, and malondialdehyde. The polyherbal tea produced no significant change in sperm count compared with the distilled-water and carbamazepine groups ($p > .05$). However, the extract significantly increased progressive sperm motility and sperm viability while reducing immotile sperm, non-progressive motility, and non-viable sperm ($p < .05$ or $p < .01$). Superoxide dismutase and catalase activities were significantly elevated

AJSTM Journal is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License

following treatment with the polyherbal tea compared with the distilled-water and carbamazepine groups ($p < .05$). The extract and sildenafil citrate also significantly reduced malondialdehyde levels ($p < .05$), whereas glutathione levels remained unchanged across the treatment groups ($p > .05$). These findings demonstrate that the aqueous polyherbal tea extract improves sperm motility and viability without affecting sperm count, potentially through enhanced testicular antioxidant activity and reduced lipid peroxidation. The study provides preliminary evidence supporting further investigation of the formulation as a potential intervention for improving male reproductive function.

Keywords: Antioxidant Activity; Male Reproductive Function; Polyherbal Tea; Sperm Motility; Sperm Viability

INTRODUCTION

Natural resources and their products are the most common sources of bioactive molecules in medicines since the development of other advanced treatments after thegenomic era (Kutama *et al.,* 2018). The World Health Organisation (WHO) describes traditional herbal medications as naturally occurring, plant-derived materials with negligible or no industrial processing which have been utilised to treat ailments within regional or local healing practices (Tilburt and Kaptchuk, 2008). Traditional systems of medicine and their formulations have been extensively used for many years in developed and developing countries, mainly due to their natural origin and minor side effects (Balunas and Kinghorn, 2005). Herbal medicine is in the mainstream of around 75–80% of the world population, primarily in the developing countries, for use in primary health care to treat diseases (Qazi and Molvi, 2016). This is mostly because of the universal belief that herbal medicines don't have any side effects, in addition to being locally available and cheap (Qazi and Molvi, 2016). The treatment of injury or disease by plants or plant material, either in the crude or processed state, is known as traditionall herbal medicine (Sadiku *et al.,* 2022). Plants have been used in a single or combined form in the treatment of diseases (Msomi and Simelane, 2017). Herb-herb combinations, also known as polyherbal therapy, have been used in Chinese medicine practice for thousands of years (Che *et al.,* 2013). Drug combination often produces a promising effect in the treatment of diseases over a single drug. The concept of drug combination has been well established in Western medicine, and remarkable success has been achieved over the decades. Though this progress is made,

scientific evidence of therapeutic benefits is still lacking in herbal medicine (Balkrishna *et al.,* 2024).

Plant Composition

Plants used include *Curcuma longa, Anthocleistadjaalonesis, Mucuna puriens, Musa paradisiaca and Thespesia garckeana.*

Plant Collection

Curcuma longa (turmeric) was purchased from Kurmi Market within Kano Municipal Local Government Area, Kano State. *Thespesia garckeana* (Gorontula) was purchased from Gombe State. *Anthocleistadjalonensis* and *Musa paradisiaca* (banana root) were gotten from Ikpoba-Okha Local Government Area in Edo State. *Mucuna pruriens* (velvet beans) was purchased from Igbanke in Orhionmwon.Local Government Area, Edo State. The plants were identified and authenticated by Dr H. Akinnibosun in the Department of Plant Biology and Biotechnology, Faculty of Life Science, University of Benin, Edo State.

Preparation of Plant Materials

Thespesia garckeana (Goron Tula) seeds were collected from the fruit. The *Curcuma longa* (turmeric) was washed in running water and chopped into smaller bits. The*sis and Mucuna pruriens (Velvet* Beans) were removed from the stalk and also washed in running water. The *Musa paradisiaca* (banana root) was air-dried and broken into smaller bits. The chopped *Curcuma longa* and *Thespesia garckeana* seeds were dehydrated using a dehydrator at 50°C, while the washed *Anthocleistadjalonensis* and *Mucuna pruriens* were air-dried. After dehydration, the dehydrated *Thespesia garckea*, *Curcuma longa*, *Mucuna pruriens*, *Musa paradisiaca* and *Anthocleistadjalonensis* were ground to powder separately using an industrial blender (KENWOOD MODEL; KCB239k). The plants were weighed separately using an analytical balance and mixed together.

Formulation of Polyherbal Tea

The polyherbal formulated tea was formulated according to Uwaya and Effiong (2024). with slight modification. The polyherbal tea was prepared using the following formulation of *Musa paradisiaca, Curcuma longa, Anthocleistadjalonensis, Mucuna pruriens and Thespesia garckeana* in the ratio of 1:1:1:0.1:0.5, respectively.

Polyherbal Tea Extraction

Formulated polyherbal tea (648 grams) was weighed into an extracting jar, and 9.75

litres of distilled water were added. The mixture was stirred using a stirrer and allowed to stand for 72 hours. The mixture was filtered using a strainer into a storage container with a lid. The filtrate was concentrated over a water bath. The extract was stored in a storage bottle in a refrigerator at 4°C prior to use.

Experimental Design

The sperm count, motility, and viability were studied according to the method of Naghdi *et al.,* 2016, with slight modification. Twenty healthy male albino rats were allotted into five groups of four rats in each group as follows:

Group 1 – Received 2 ml/kg of distilled water.

Group 2 – Received 20 mg/kg of carbamazepine (standard drug).

Group 3 received 5 mg/kg of sildenafil (standard drug).

Group 4 – Received 10 mg/kg aqueous extract of polyherbal-formulated tea.

Group 5 – Received 20 mg/kg aqueous extract of polyherbal-formulated tea.

The control group, standard group, and extracts group were administered orally for 21 days.

Epididymal Sperm Preparation

After 21 days, the animals were sacrificed using chloroform as anaesthesia, after which the testes, epididymis, and seminal vesicles were removed, isolated, and transferred into a Petri dish containing 1 ml of phosphate buffer after ensuring the adherent tissue was cleared from the cordial epididymis. The caudal epididymis was cut open to release the sperm in it and was left for 15-30 minutes.

Sperm Count Assessment

The sperm count was evaluated by diluting one ml of sperm suspension with one ml of sperm dilution fluid. Ten microlitres of the mixture were transferred into a hemocytometer, and the sperm count was assessed per 250 tiny squares of the hemocytometer (Naghdi *et al.,* 2016).

Sperm Motility Assessment

Two drops of the sperm were dropped on a clean slide with the aid of a Pasteur pipette. The slides were covered with a cover slip and viewed under a light microscope at x40 magnification. The progressive motility, nonprogressive motility and immotile were observed, counted and recorded (Naghdi *et al.,* 2016).

Sperm Viability Assessment

To determine the percentage of viable sperm, 7 μL of trypan blue was mixed with 20 μL of sperm suspension. The percentage of live (unstained) and dead (blue stained) spermatozoa was measured using a light microscope at x400 magnification (Naghdi *et al.,* 2016).

Histological Assessment

The tissue specimen was dehydrated in alcohol, processed in xylene and then embedded in paraffin wax before sectioning with a microtome. Sections (5µm thick) of the tissues were stained with haematoxylin and eosin. All sections were viewed under a Leica light microscope (Model DM500).

RESULTS

The effect of polyherbal formulated tea (*Mucuna pruriens*, *Curcuma longa, Anthocleista djanalonsis, Musa paradisaca and Thespesia garckeana)* on sperm count, viability and motility




Figure 1: The effect of polyherbal formulated tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on sperm count,

The polyherbal formulated tea at 10 and 20 mg/kg has no effect on the sperm count when compared with normal control and Carbamazepine ($p > 0.05$). While Sildenafil Citrate showed a significant increase in sperm count when compared with normal,

Carbamazepine and the polyherbal formulated tea at 10 and 20 mg/kg (**p < 0.01). Values were presented as mean ± S.E.M. n=4




Figure 2: The effect of polyherbal formulated tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Sperm Progressive Motility.

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg and Sildenafil Citrate significantly increased sperm progressive motility when compared with water control and Carbamazepine (**p<0.01; *p<0.05). Values were presented as mean ± S.E.M. n=4




Figure 3: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Non Progressive Sperm Motility.

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg and sildenafil citrate significantly reduced non-progressive sperm motility when compared to carbamazepine (**p < 0.01). Values were presented as mean ± S.E.M. n=4




Figure 4: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on immotile sperm.

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg significantly reduced immotile sperm when compared with water control and carbamazepine (**p < 0.01). Values were presented as mean ± S.E.M. n=4




Figure 5: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Viable Sperm.

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg and Sildenafil Citrate significantly increased the viable sperm when compared with water control and Carbamazepine (**p<0.01; *p<0.05). Values were presented as the mean ± S.E.M. n = 4.




Figure 6: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Non Viable

Sperm.

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg significantly reduced the non-viable sperm when compared with carbamazepine and normal control (*p<0.05; *p<0.05). Values were presented as mean ± S.E.M. n=4




Figure 7: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Superoxide dismutase (SOD).

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg significantly increased the superoxide dismutase (SOD) when compared with the normal control (*p<0.05). Values were presented as the mean ± S.E.M. n = 4.




Figure 8: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Catalase (CAT).

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg significantly increased the Catalase (CAT) when compared with water control. (**p<0.01). Values were presented as mean ± S.E.M. n=4.




Figure 9: The effect of polyherbal formulated tea tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana*) on Malondialdehyde (MDA).

The aqueous extract of polyherbal formulated tea at 10 and 20 mg/kg and sildenafil citrate significantly reduced malondialdehyde (MDA) when compared with normal control and carbamazepine (**p<0.01; *p<0.05). Values were presented as the mean ± S.E.M. n=4.





Figure 10: tea (*Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca and Thespesia garckeana* on Glutathione (GSH).

The aqueous extracts of polyherbal-formulated tea at 10 and 20 mg/kg, carbamazepine, and sildenafil citrate have no significant effect on glutathione (GSH) when compared with water control (p > 0.05). Values were presented as the mean ± S.E.M. n = 4.






Plate A-Control: A section of the testis shows oval-shaped seminiferous tubules (thick arrow) containing Sertoli cells and sperm cells at different stages of maturation. The tubules are surrounded by a thin membrane with the presence of Leydig cells in the interstitium (thin arrow). Features are in keeping with normal testicular tissue.

Plate B-Sildenafil Citrate 5 5mg/kg: Section of the testis shows oval-shaped seminiferous tubules (thick arrow) containing Sertoli cells and sperm cells at different stages of maturation. The tubules are surrounded by a thin membrane with the presence of Leydig cells in the interstitium (thin arrow). Features are in keeping with normal testicular tissue.

Plate C – Carbamazepine 20 mg/kg: A section of the testis shows oval-shaped seminiferous tubules (thick arrow) containing Sertoli cells and sperm cells at different stages of maturation. The tubules are surrounded by a thin membrane with the presence of Leydig cells in the interstitium (thin arrow). Features are in keeping with normal testicular tissue.

Plate D - 10 mg/kg: A section of the testis shows oval-shaped seminiferous tubules (thick arrow) containing Sertoli cells and sperm cells at different stages of maturation. The tubules are surrounded by a thin membrane with the presence of Leydig cells in the interstitium (thin arrow). Features are in keeping with normal testicular tissue.

Plate E - 20 mg/kg: A section of the testis shows oval-shaped seminiferous tubules (thick arrow) containing Sertoli cells and sperm cells at different stages of maturation. A

thin membrane surrounds the tubules, and the interstitium contains Leydig cells (thin arrow). Features are in keeping with normal testicular tissue**.**

DISCUSSION

In traditional medicine, herbs such as *Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca,* and *Thespesia garckeana* are used to treat various ailments such as malaria, diabetes, neurodegenerative disorders, and gastrointestinal issues (Chinaza *et al.,* 2019; Enwerem *et al.,* 2019; Gondwe *et al.,* 2018; Balunas and Kinghorn, 2005). Researchers have also found them useful in treating infertility by enhancing the motility and viability of sperm. Infertility refers to the inability to conceive after at least one year of unprotected intercourse (Larsen, 2005). In this study, the teas made from *Anthocleista djalonesis, Curcuma longa, Mucuna pruriens, Musa paradisiaca*, and *Thespesia garckeana* at doses of 10 mg/kg and 20 mg/kg did not change the number of sperm counts, while the standard treatment (Sildenafil Citrate 5 mg/kg) increased the sperm count, as shown in Figure 1. Polyherbal tea at 10 and 20 mg/kg and sildenafil citrate significantly increased viable sperm (Figure 5) and reduced non-progressive sperm as well as non-viable sperm (Figures 2 and 6). The ability of a plant to increase sperm viability and progressive sperm motility suggests that the plant possesses properties linked to promoting male fertility (Khaki *et al.,* 2009). The polyherbal-formulated tea's ability to increase sperm viability and progressive sperm motility suggests that the tea possesses properties linked to promoting male fertility. Carbamazepine used as a negative control increased non-progressive sperm and immotile sperm, as seen in figures 3 and 4 (Asadi-Pooya *et al.,* 2015). Carbamazepine (CBZ) is one of the main drugs used to treat epilepsy and neuropathic pain (Boon *et al.,* 2021). Researchers recognise that carbamazepine (CBZ) and its metabolites induce oxidative stress and reactive oxygen species (ROS) (Mittag *et al.,* 2016). Carbamazepine (CBZ) administration reduces sperm concentration, testicular heaviness, testosterone, FSH, and LH hormones (Akorede *et al.,* 2020; Boon *et al.,* 2021; Asadi-Pooya *et al.,* 2015; Kawata *et al.,* 2001). Sildenafil citrate acts by increasing blood flow to the penis during sexual stimulation (Loran *et al.,* 2009). A study shows that *M. pruriens* improved sperm concentration and motility (Ahmad *et al*., 2008). The ability of *Mucuna pruriens t*o boost sperm concentration and movement is linked to its active ingredients, like alkaloids, coumarins, flavonoids, and alkylamines, which may help enhance the antioxidant levels in men's fertility (Mishra *et al.,* 2024; Ahmad *et al.,*

2008*)*. The phytochemicals in *Thespesia garckeana* have been studied for their potential impact on male fertility, and the plant contains flavonoids, alkaloids, and other compounds that may contribute to improved male reproductive health, including increased testosterone levels. These phytochemicals are believed to play a role in modulating reproductive hormones and supporting healthy testicular function **(**Bukar *et al*., 2021). *Anthocleista djalonensis* also contains flavonoids, alkaloids, phenol and other compounds that may contribute to improved male reproductive health (Popoola, 2020). The phytochemicals in *velvet bean (Mucuna pruriens), Goron Tula (Thespesia garckeana), and Anthocleista djalonensis* might be the reason for the improvement in sperm quality found in this study. Researchers have found that the Velvet bean (*Mucuna pruriens*), the Goron Tula (*Thespesia garckeana*), and the *Anthocleista djalonensis* can help with fertility (Okeke *et al.,* 2019; Maroyi, 2017; Karitha and Thangamam, 2014). McLeod was the first to suggest that oxidative stress may play a role in the aetiology of defective sperm functionality (Aitken *et al.,* 2014). A decrease in seminal plasma antioxidant capacity (TAC) could play a crucial role in the pathogenesis of reduced sperm function. Therefore, we can use the total antioxidant capacity (TAC) as a specific biomarker to gauge oxidative stress in sperm (Pahune *et al*., 2013). At 10 and 20 mg/kg, the aqueous extract of polyherbal-formulated tea significantly increased superoxide dismutase (SOD). It is widely accepted that SOD is the body's first defence against the harmful effects of $\bullet O_2-$ because it helps convert two $\bullet O_2-$ molecules into hydrogen peroxide and O_2, which reduces the amount of $\bullet O_2-$ available, and it plays a significant role in maintaining sperm motility (Öner-İyidoğan *et al*., 2003). Low and diminished SOD activity has been associated with a significant risk of oxidative stress, resulting in diseases such as hypertension, hypercholesterolaemia, atherosclerosis, diabetes, heart failure, stroke, and other cardiovascular diseases (Rosa *et al.,* 2021; Fukai and Ushio-Fukai, 2011). An increase of SOD level by the herbal tea in this study shows that the polyherbal tea may neutralise free radicals that may slow down sperm motility and improve sperm health. In this study, the aqueous extract of polyherbal-formulated tea at 10 and 20 mg/kg significantly increased catalase (CAT). Catalase (CAT) treats many oxidative stress illnesses (Nandi *et al.,* 2019). It is an important enzyme in seminal plasma that aids sperm's antioxidant (Nandi *et al.,* 2019). It contributes to maintaining normal reactive oxygen species (ROS) levels while also protecting sperm from potentially harmful ROS levels. Catalase (CAT) reduces H_2O_2, which is an essential sensor in redox metabolism, to water. H_2O_2 levels are critical to oxidative stress. At 10 and 20 mg/kg, the aqueous extract of polyherbal-formulated tea and

sildenafil citrate significantly reduced malondialdehyde (MDA). Sperm cell membranes are vulnerable to peroxidative damage; malondialdehyde (MDA) serves as an index of lipid peroxidation and a marker of oxidative stress, which may be used as a diagnostic tool for analysing infertility in asthenozoospermic patients (Tavilani*et al.,* 2005). Researchers have found that higher levels of malondialdehyde in the blood are linked to lower sperm health, including less viable sperm, reduced movement, abnormal shape, and lower sperm count. The ability of the tea to lower malondialdehyde (MDA) levels indicates that the polyherbal tea promotes sperm health. According to this research, the aqueous extract of polyherbal-formulated tea at 10 and 20 mg/kg has no significant effect on glutathione (GSH). The testicular histopathology has been, for years, one of the crucial investigations in the diagnosis and management of male infertility (Dohle *et al.,* 2012). One of the major challenges with conventional TESE (cTESE) was finding indicators of the likelihood of recovering spermatozoa. Occasionally, circulating hormonal markers, testicular ultrasound, and testicular histopathology have been used (Guler *et al.,* 2016). Histological slides of the testes showed that they were normal (Plates A–E). Malondialdehyde, catalase, and superoxide dismutase significantly influence the potential for sperm fertilisation and male infertility (Tavilani*e et al.,* 2005). A polyherbal tea containing *Anthocleista djalonensis, Curcuma longa, Mucuna pruriens, Musa paradisiaca, and Thespesia garckeana* may boost male fertility, according to this study.

CONCLUSION

The polyherbal-formulated tea has no effect on sperm count, but it does increase sperm motility and viability. It also possesses antioxidant activity.

Acknowledgement:

We are quite grateful to Mr. A. Barnabas and Mr. Ediri Vaikosen, who work in the laboratory of the Department of Science Laboratory Technology at the Faculty of Life Sciences, for their essential assistance with our research.

Conflict of Interests

There is no conflict between the study's conduct and the authors' interests.

Authors' Contributions

DOU participated in the manuscript's drafting, as well as the study's design and supervision. CCU and JNO actively participated in both the experimentation and the drafting of the manuscript. Before submitting the work, all authors reviewed it and gave their approval.

REFERENCES

Adebayo, A. A. Ogunboye, & Olamide, O. O. Crown. (2022). *Anthocleista djalonensis*: A review of its ethnobotanical, phytochemical and pharmacological potentials. *Journal of Clinical and Medical Reviews, 1*(1). https://www.mediresonline.org/anthocleista-djalonensis-a-review-of-its-ethnobotanical-phytochemical-and-pharmacological-potentials.html

Adeeko, A. O., Bakare, A. A., Adeniji, A. O., & Nwachukwu, N. (2016). Antioxidative and hypoglycemic activities of the African herb *Azanza garckeana* in alloxan-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine, 6*(1), 54–60.

Adenowo, A. F., Salisu, F. T., Akinsanya, M. A., & Kazeem, M. I. (2022). Nutraceutical potentials of *Azanza garckeana* (snot apple): A review. *Current Nutraceuticals, 3*(2). https://www.eurekaselect.com/public/article/121738

Agarwal, A., Deepinder, F., Sharma, R. K., Ranga, G., & Li, J. (2008). Effect of cell phone usage on semen analysis in men attending infertility clinic: An observational study. *Fertility and Sterility, 89*(1), 124–128. https://doi.org/10.1016/j.fertnstert.2007.01.166

Agarwal, A., Mulgund, A., Hamada, A., & Chyatte, M. R. (2015). A unique view on male infertility around the globe. *Reproductive Biology and Endocrinology, 13*, Article 37. https://doi.org/10.1186/s12958-015-0032-1

Agrawal, A., Panis, B., & Swennen, R. (2002). Effect of pretreatment conditions and genotype on cryopreservation of in vitro cultures of banana meristems. *Proceedings of the Third International Symposium on Molecular and Cellular Biology of Banana, 49*, 9–11.

Agrawal, A., Swennen, R., & Panis, B. (2004). A comparison of four methods for cryopreservation of meristems in banana (*Musa* spp.). *CryoLetters, 25*, 101–110.

Ahmad, M. K., Mahdi, A. A., Shukla, K. K., Islam, N., Jaiswar, S. P., & Ahmad, S. (2008). Effect of *Mucuna pruriens* on semen profile and biochemical parameters in seminal plasma of infertile men. *Fertility and Sterility, 90*(3), 627–635. https://doi.org/10.1016/j.fertnstert.2007.07.1314

Aitken, R. J., Smith, T. B., Jobling, M. S., Baker, M. A., & De Iuliis, G. N. (2014). Oxidative stress and male reproductive health. *Asian Journal of Andrology, 16*(1), 31–38. https://doi.org/10.4103/1008-682X.122203

Ajayi, F. A., Ogunleye, O. O., Adelowo, O. O., & Adenuga, G. A. (2020). The chemical composition, antioxidant, anti-inflammatory, and antifertility potentials of *Anthocleista djalonensis* on reproductive hormones in male albino rats. *Clinical Phytoscience, 6*, Article 58.

Akorede, G. J., Ambali, S. F., Olatunji, A. O., Shittu, M., Aremu, A., Basiru, A., Mikai'l, G. H., Khalid, T. B., Akeem, O. A., & Saliu, A. A. (2020). Protective effect of vitamin

C on chronic carbamazepine-induced reproductive toxicity in male Wistar rats. *Toxicology Reports, 7*, 269–276.

Akram, M., Shahab-Uddin, Usmanghani, K., Hannan, A., Mohiuddin, E., & Asif, M. (2010). *Curcuma longa* and curcumin: A review article. *Romanian Journal of Biology—Plant Biology, 55*(2), 65–70.

Anyanwu, G. O., Ur-Rehman, N., Onyeneke, C. E., & Rauf, K. (2015). Medicinal plants of the genus *Anthocleista*—A review of their ethnobotany, phytochemistry and pharmacology. *Journal of Ethnopharmacology, 175*, 648–667. https://doi.org/10.1016/j.jep.2015.09.032

Asadi-Pooya, A. A., Farazdaghi, M., & Ashjazadeh, N. (2015). Effects of carbamazepine on semen parameters in men with newly diagnosed epilepsy. *Iranian Journal of Neurology, 14*(3), 168–170.

Aw, T. Y., Wierzbicka, G., & Jones, D. P. (1991). Oral glutathione increases tissue glutathione in vivo. *Chemico-Biological Interactions, 80*(1), 89–97. https://doi.org/10.1016/0009-2797(91)90033-4

Ayman, E. L., Hassan, S. M., Mohamed, A. M. M., & Mohammed, H. M. A. (2019). Tumeric or *Curcuma longa* Linn. In *Nonvitamin and nonmineral nutritional supplements* (pp. 447–453). Academic Press. https://doi.org/10.1016/B978-0-12-812491-8.00059-X

Ballatori, N., Krance, S. M., Notenboom, S., Shi, S., Tieu, K., & Hammond, C. L. (2009). Glutathione dysregulation and the etiology and progression of human diseases. *Biological Chemistry, 390*(3), 191–214. https://doi.org/10.1515/BC.2009.033

Balunas, M. J., & Kinghorn, A. D. (2005). Drug discovery from medicinal plants. *Life Sciences, 78*(5), 431–441. https://doi.org/10.1016/j.lfs.2005.09.012

Banerjee, N., & De Langhe, E. (1985). A tissue culture technique for rapid clonal propagation and storage under minimal growth conditions of *Musa* (banana and plantain). *Plant Cell Reports, 4*(6), 351–354. https://doi.org/10.1007/BF00269897

Ben Abdallah, F., Dammak, I., Attia, H., Hentati, B., & Ammar-Keskes, L. (2009). Lipid peroxidation and antioxidant enzyme activities in infertile men: Correlation with semen parameters. *Journal of Clinical Laboratory Analysis, 23*(2), 99–104.

Bioltif, Y. E., Edward, N. B., & Tyeng, T. D. (2020). A chemical overview of *Azanza garckeana*. *Biology, Medicine, & Natural Product Chemistry, 9*(2), 91–95. https://doi.org/10.14421/biomedich.2020.92.91-95

Boon, P., Ferrao Santos, S., Jansen, A. C., Lagae, L., Legros, B., & Weckhuysen, S. (2021). Recommendations for the treatment of epilepsy in adult and pediatric patients in Belgium: 2020 update. *Acta Neurologica Belgica, 121*(1), 241–257. https://doi.org/10.1007/s13760-020-01488-y

Bravard, A., Sabatier, L., Hoffschir, F., Ricoul, M., Luccioni, C., & Dutrillaux, B. (1992). SOD2: A new type of tumor-suppressor gene? *International Journal of Cancer, 51*(3), 476–480. https://doi.org/10.1002/ijc.2910510323

Buckman, L. (1998). The utilization of velvet beans in various regions. *Journal of Agricultural Studies, 12*(3), 45–58.

Bukata, B., Bukar, F. E., & Sunday, Y. S. (2021). Methanol extract of *Azanza garckeana* fruit pulps protects against formalin-induced reproductive toxicity in adult albino male

mice. *Journal of Advances in Medical and Pharmaceutical Sciences, 23*(8), 38–49. https://doi.org/10.9734/jamps/2021/v23i830253

Che, C.-T., Wang, Z. J., Chow, M. S. S., & Lam, C. W. K. (2013). Herb–herb combination for therapeutic enhancement and advancement: Theory, practice and future perspectives. *Molecules, 18*(5), 5125–5141. https://doi.org/10.3390/molecules18055125

Cho, C. L., Esteves, S. C., & Agarwal, A. (2018). Novel insights into the pathophysiology of varicocele and its association with reactive oxygen species and sperm DNA fragmentation. *Asian Journal of Andrology, 20*(4), 294–301. https://doi.org/10.4103/aja.aja_107_17

Choudhury, D., Bhattacharyya, S., Bose, S., & Gupta, P. (2013). Curcumin suppresses nuclear factor-κB-mediated inflammation in injured spinal cord of rats. *Journal of Surgical Research, 183*(1), 163–175.

Corona, G., Rastrelli, G., Monami, M., Saad, F., Luconi, M., Lucchese, M., Facchiano, E., Sforza, A., Forti, G., & Mannucci, E. (2013). Body mass index regulates hypogonadism-associated cardiovascular risk: Results from a cohort of subjects with erectile dysfunction. *The Journal of Sexual Medicine, 10*(12), 3102–3109.

Corvo, M. L., Jorge, J. C., van't Hof, R., Cruz, M. E. M., Crommelin, D. J. A., & Storm, G. (2002). Superoxide dismutase entrapped in long-circulating liposomes: Formulation design and therapeutic activity in rat adjuvant arthritis. *Biochimica et Biophysica Acta—Biomembranes, 1564*(1), 227–236.

Deisseroth, A., & Dounce, A. L. (1970). Catalase: Physical and chemical properties, mechanism of catalysis, and physiological role. *Physiological Reviews, 50*(3), 319–375. https://doi.org/10.1152/physrev.1970.50.3.319

Deokar, G., Kakulte, H., & Kshirsagar, S. (2016). Phytochemistry and pharmacological activity of *Mucuna pruriens*: A review. *Pharmaceutical and Biological Evaluations, 3*(1), 50–59.

Dikko, Y. J., Khan, M. E., Tor-Anyiin, T. A., Anyam, J. V., & Linus, U. A. (2016). In vitro antimicrobial activity of fruit pulp extracts of *Azanza garckeana* (F. Hoffm.) Exell & Hillc. and isolation of one of its active principles, betulinic acid. *British Journal of Pharmaceutical Research, 14*(1), 1–10. https://doi.org/10.9734/BJPR/2016/30152

Dissanayaka, D., Udumann, S. S., Nuwarapaksha, T. D., & Atapattu, A. J. (2024). Harnessing the potential of *Mucuna* cover cropping: A comprehensive review of its agronomic and environmental benefits. *Circular Agricultural Systems, 4*(1), 1–9.

Dohle, G. R., Elzanaty, S., & van Casteren, N. J. (2012). Testicular biopsy: Clinical practice and interpretation. *Asian Journal of Andrology, 14*(1), 88–93. https://doi.org/10.1038/aja.2011.57

Drosdzol, A., & Skrzypulec, V. (2009). Evaluation of marital and sexual interactions of Polish infertile couples. *The Journal of Sexual Medicine, 6*(12), 3335–3346.

Duty, S. M., Silva, M. J., Barr, D. B., Brock, J. W., Ryan, L., Chen, Z., Herrick, R. F., Christiani, D. C., & Hauser, R. (2003). Phthalate exposure and human semen parameters. *Epidemiology, 14*(3), 269–277.

Eilittä, M., Mureithi, J., & Derpsch, R. (2005). Cover crops in smallholder maize production systems of East Africa. In A. Bationo (Ed.), *Managing nutrient cycles to sustain soil fertility in Sub-Saharan Africa*. Academy Science Publishers.

Eldad, A., Ben Meir, P., Breiterman, S., Chaouat, M., Shafran, A., & Ben-Bassat, H. (1998). Superoxide dismutase for mustard gas burns. *Burns, 24*(2), 114–119.

Elkhoudri, N., Baali, A., Habbal, R., Benmoussa, A., & Amarouch, H. (2018). The effect of *Musa paradisiaca* flowers on the reproductive system of rats. *Journal of Herbal Medicine, 11*, 43–48.

Esteves, S. C., & Agarwal, A. (2013). Novel concepts in male infertility. *International Brazilian Journal of Urology, 39*(1), 12–20. https://doi.org/10.1590/S1677-5538.IBJU.2013.01.03

Esteves, S. C., Roque, M., & Bedoschi, G. (2020). Systematic review of clinical practice guidelines for failed in vitro fertilization cycles. *JBRA Assisted Reproduction, 24*(4), 480–492.

Forman, H. J., Zhang, H., & Rinna, A. (2009). Glutathione: Overview of its protective roles, measurement, and biosynthesis. *Molecular Aspects of Medicine, 30*(1–2), 1–12. https://doi.org/10.1016/j.mam.2008.08.006

Fukai, T., & Ushio-Fukai, M. (2011). Superoxide dismutases: Role in redox signaling, vascular function, and diseases. *Antioxidants & Redox Signaling, 15*(6), 1583–1606. https://doi.org/10.1089/ars.2011.3999

Fuloria, S., Mehta, J., Chandel, A., Sekar, M., Rani, N. N. I. M., Begum, M. Y., Subramaniyan, V., Chidambaram, K., Thangavelu, L., Nordin, R., & Wu, Y. S. (2022). A comprehensive review on the therapeutic potential of *Curcuma longa* Linn. in relation to its major active constituent curcumin. *Frontiers in Pharmacology, 13*, 820806. https://doi.org/10.3389/fphar.2022.820806

Gameiro, S., Boivin, J., Peronace, L., Verhaak, C. M., & Wischmann, T. (2012). Why do patients discontinue fertility treatment? A systematic review of reasons and predictors of discontinuation in fertility treatment. *Human Reproduction Update, 18*(6), 652–669. https://doi.org/10.1093/humupd/dms031

Gbadamosi, I. T., & Erinoso, S. M. (2016). A review of twenty ethnobotanicals used in the management of breast cancer in Abeokuta, Ogun State, Nigeria. *African Journal of Pharmacy and Pharmacology, 10*(27), 546–564.

George, S. K., Rajesh, R., Kumar, S. S., Sulekha, B., & Balaram, P. (2008). A polyherbal Ayurvedic drug, Indukantha Ghritha, as an adjuvant to cancer chemotherapy via immunomodulation. *Immunobiology, 213*(8), 641–649.

Giustarini, D., Dalle-Donne, I., Milzani, A., & Rossi, R. (2019). Red blood cells as a physiological source of glutathione for extracellular fluids. *Blood Cells, Molecules, and Diseases, 76*, 25–29.

Greil, A. L., Slauson-Blevins, K., & McQuillan, J. (2010). The experience of infertility: A review of recent literature. *Sociology of Health & Illness, 32*(1), 140–162.

Güler, I., Erdem, M., & Erdem, A. (2016). Impact of testicular histopathology as a predictor of sperm retrieval and pregnancy outcome in patients with nonobstructive azoospermia: Correlation with clinical and hormonal factors. *Andrologia, 48*, 765–773.

Hinton, L., Miller, T., Wertz, S., & Reilly, N. (2019). Relationships between coping, coping assistance, and depression among couples undergoing infertility treatment. *Journal of Behavioral Medicine, 42*(2), 314–325.

Hossain, M. K., Al-Raqmi, K. A. S., Al-Mijizy, Z. H., Weli, A. M., & Al-Riyami, Q. (2011). Study of total phenol and flavonoid contents and phytochemical screening of various crude leaf extracts of locally grown *Thymus vulgaris*. *Asian Pacific Journal of Tropical Biomedicine, 1*(5), 395–398.

Houston, M. C. (2011). The role of potassium in hypertension and cardiovascular disease. *The Journal of Clinical Hypertension, 13*(6), 414–422.

Hsieh, Y.-Y., Chang, C.-C., & Lin, C.-S. (2006). Seminal malondialdehyde concentration but not glutathione peroxidase activity is negatively correlated with sperm concentration and motility. *International Journal of Biological Sciences, 2*(1), 23–29. https://doi.org/10.7150/ijbs.2.23

Ivancich, A., Jouve, H. M., Sartor, B., & Gaillard, J. (1997). EPR investigation of compound I in *Proteus mirabilis* and bovine liver catalases: Formation of porphyrin and tyrosyl radical intermediates. *Biochemistry, 36*(31), 9356–9364.

Iweala, E. J., Uche, M. E., Dike, E. D., Etumnu, L. R., Dokunmu, T. M., Oluwapelumi, A. E., Okoro, B. C., Dania, O. E., Adebayo, A. H., & Ugbogu, E. A. (2023). *Curcuma longa* (turmeric): Ethnomedicinal uses, phytochemistry, pharmacological activities and toxicity profiles—A review. *Pharmacological Research—Modern Chinese Medicine, 6*, 100222. https://doi.org/10.1016/j.prmcm.2023.100222

Jones, D. P., & Liang, Y. (1998). Glutathione in plasma and blood: Relationships to biology and pathology. *Methods in Enzymology, 300*, 3–8.

Jyotirmayee, B., & Mahalik, G. (2022). A review on selected pharmacological activities of *Curcuma longa* L. *International Journal of Food Properties, 25*(1), 1377–1398.

Kajaria, D. K., Gangwar, M., Sharma, A. K., Nath, G., & Bhusan, Y. (2011). Comparative evaluation of phenol and flavonoid content of polyherbal drugs. *Pharmacologyonline, 3*, 1365–1373.

Kangralkar, V. A., Patil, S. D., & Bandivadekar, R. M. (2010). Oxidative stress and diabetes: A review. *International Journal of Applied Pharmaceutics, 1*(1), 38–45.

Kapusta, A., Kuczyńska, B., & Puppel, K. (2018). Relationship between the degree of antioxidant protection and the level of malondialdehyde in high-performance Polish Holstein–Friesian cows at peak lactation. *PLOS ONE, 13*(3), e0193512. https://doi.org/10.1371/journal.pone.0193512

Karlowicz-Bodalska, K., Han, S. T., Freier, J., Smoleński, M., & Bodalska, A. (2017). *Curcuma longa* as a medicinal herb in the treatment of diabetic complications. *Acta Poloniae Pharmaceutica—Drug Research, 74*(2), 605–610.

Kavitha, C., & Thangamani, C. (2014). Amazing bean *Mucuna pruriens*: A comprehensive review. *Journal of Medicinal Plants Research, 8*, 138–143.

Kawata, Y., Okada, M., Murakami, T., Kamata, A., Zhu, G., & Kaneko, S. (2001). Pharmacological discrimination between effects of carbamazepine on hippocampal basal, Ca^{2+}- and K^{+}-evoked serotonin release. *British Journal of Pharmacology, 133*(4), 557–567.

Khera, M. (2018). *Male infertility: A clinical guide*. Springer.

Kocaadam, B., & Şanlier, N. (2017). Curcumin, an active component of turmeric (*Curcuma longa*), and its effects on health. *Critical Reviews in Food Science and Nutrition, 57*(13), 2889–2895. https://doi.org/10.1080/10408398.2015.1077195

Kohn, T. P., Kohn, J. R., & Pastuszak, A. W. (2019). Varicocele: Practical guidelines for evaluation and treatment. *Seminars in Interventional Radiology, 36*(3), 212–221. https://doi.org/10.1055/s-0039-1697909

Kunwar, R. M., Shrestha, K. P., & Bussmann, R. W. (2010). Traditional herbal medicine in far-west Nepal: A pharmacological appraisal. *Journal of Ethnobiology and Ethnomedicine, 6*, Article 35. https://doi.org/10.1186/1746-4269-6-35

Kutama, R. M., Abdulkadir, S., Kwalli, S. A., & Chiroma, G. (2018). Phytochemical compositions in some Nigerian medicinal plants and their pharmacological properties: A review. *International Journal of Anesthesia and Clinical Medicine, 6*(1), 15–25.

Lal, J. (2012). Turmeric, curcumin and our life: A review. *Bulletin of Environment, Pharmacology and Life Sciences, 1*(7), 11–17.

Lampariello, L. R., Cortelazzo, A., Guerranti, R., Sticozzi, C., & Valacchi, G. (2012). The magic velvet bean of *Mucuna pruriens*. *Journal of Traditional and Complementary Medicine, 2*(4), 331–339. https://doi.org/10.1016/S2225-4110(16)30119-5

Larsen, U. (2005). Research on infertility: Which definition should we use? *Fertility and Sterility, 83*(4), 846–852. https://doi.org/10.1016/j.fertnstert.2004.11.033

Lawal, B., Sani, S., Onikanni, A. S., Ibrahim, Y. O., Agboola, A. R., Lukman, H. Y., Olawale, F., Jigam, A. A., Batiha, G. E.-S., Babalola, S. B., Mostafa-Hedeab, G., Lima, C. M. G., Wu, A. T.-H., Huang, H.-S., & Conte-Junior, C. A. (2022). Preclinical anti-inflammatory and antioxidant effects of *Azanza garckeana* in STZ-induced glycemic-impaired rats, and pharmacoinformatics of its major phytoconstituents. *Biomedicine & Pharmacotherapy, 152*, 113196. https://doi.org/10.1016/j.biopha.2022.113196

Leonti, M., & Casu, L. (2013). Traditional medicines and globalization: Current and future perspectives in ethnopharmacology. *Frontiers in Pharmacology, 4*, Article 92. https://doi.org/10.3389/fphar.2013.00092

Liu, D., Li, L., & Augustus, L. (2001). Prostaglandin release by spinal cord injury mediates production of hydroxyl radical, malondialdehyde and cell death: A site of the neuroprotective action of methylprednisolone. *Journal of Neurochemistry, 77*(4), 1036–1047.

Lotti, F., & Maggi, M. (2018). Hormonal treatment for the management of male infertility. *Asian Journal of Andrology, 20*(2), 120–123. https://doi.org/10.4103/aja.aja_52_17

Lu, S. C. (2013). Glutathione synthesis. *Biochimica et Biophysica Acta—General Subjects, 1830*(5), 3143–3153. https://doi.org/10.1016/j.bbagen.2012.09.008

Maroyi, A. (2017). *Azanza garckeana* fruit tree: Phytochemistry, pharmacology, nutritional and primary healthcare applications as herbal medicine—A review. *Research Journal of Medicinal Plant, 11*, 115–123. https://doi.org/10.3923/rjmp.2017.115.123

McCord, J. M., & Fridovich, I. (1969). Superoxide dismutase: An enzymic function for erythrocuprein. *Journal of Biological Chemistry, 244*(22), 6049–6055.

Menezes, E. W., Tadini, C. C., Tribess, T. B., Zuleta, A., Binaghi, J., Pak, N., Vera, G., Dan, M. C. T., Bertolini, A. C., & Cordenunsi, B. R. (2006). Chemical composition and nutritional value of unripe banana flour (*Musa acuminata*, var. Nanicão). *Plant Foods for Human Nutrition, 61*(2), 63–68. https://doi.org/10.1007/s11130-006-0028-4

Michael, K. G., Onyia, L. U., & Jidauna, S. B. (2015). Evaluation of phytochemicals in *Azanza garckeana* (Goron Tula) seed. *IOSR Journal of Agriculture and Veterinary Science, 8*(5), 71–74.

Milewicz, A., Gejdel, E., Sworen, H., Sienkiewicz, K., Jedrzejak, J., & Teucher, T. (1993). *Vitex agnus castus* extract in the treatment of luteal phase defects due to latent hyperprolactinemia. *Arzneimittel-Forschung, 43*(7), 752–756.

Mishra, R., Nikam, A., Hiwarkar, J., et al. (2024). Flavonoids as potential therapeutics in male reproductive disorders. *Future Journal of Pharmaceutical Sciences, 10*, Article 100.

Mittag, N., Meister, S., Berg, A., & Walther, U. (2016). A case report of a carbamazepine overdose with focus on pharmacokinetic aspects. *Pharmacopsychiatry, 49*(2), 76–78.

Mokbel, M. S., & Hashinaga, F. (2005). Antibacterial and antioxidant activities of banana (*Musa*, AAA cv. Cavendish) fruit peel. *American Journal of Biochemistry and Biotechnology, 1*(3), 125–131. https://doi.org/10.3844/ajbbsp.2005.125.131

Mwove, J. K. (2021). Potential for development of novel food products from *Azanza garckeana* tree fruit: A review. *International Journal of Food Studies, 10*, SI121–SI132. https://doi.org/10.7455/ijfs/10.SI.2021.a10

Nandi, A., Shah, A., & Wilkinson, H. (2018). Ovarian hyperstimulation syndrome: A clinical review. *Journal of Human Reproductive Sciences, 11*(4), 309–318.

Nandi, A., Yan, L.-J., Jana, C. K., & Das, N. (2019). Role of catalase in oxidative stress- and age-associated degenerative diseases. *Oxidative Medicine and Cellular Longevity, 2019*, 9613090. https://doi.org/10.1155/2019/9613090

Ngo, L. T., Okogun, J. I., & Folk, W. R. (2013). 21st century natural product research and drug development and traditional medicines. *Natural Product Reports, 30*(4), 584–592. https://doi.org/10.1039/C3NP20120A

Nieman, D. C., Gillitt, N. D., Henson, D. A., Sha, W., Shanely, R. A., Knab, A. M., Cialdella-Kam, L., & Jin, F. (2012). Bananas as an energy source during exercise: A metabolomics approach. *PLOS ONE, 7*(5), e37479. https://doi.org/10.1371/journal.pone.0037479

Nunomura, A., Moreira, P. I., Castellani, R. J., Lee, H.-G., Zhu, X., Smith, M. A., & Perry, G. (2012). Oxidative damage to RNA in aging and neurodegenerative disorders. *Neurotoxicity Research, 22*, 231–248.

O'Flynn O'Brien, K. L., Varghese, A. C., & Agarwal, A. (2010). The genetic causes of male factor infertility: A review. *Fertility and Sterility, 93*(1), 1–12. https://doi.org/10.1016/j.fertnstert.2009.10.045

Okeke, U. B., Igbinaduwa, P. O., & Ozolua, R. I. (2019). Effects of *Anthocleista djalonensis* (Chev.) methanol root-bark extract on some fertility parameters in male rats. *Nigerian Journal of Pharmaceutical Research, 15*(2), 187–192.

Öner-İyidoğan, Y., Genç, S., Koçak, H., & Akkuş, E. (2003). The effects of superoxide dismutase activity and total antioxidant status in seminal plasma on male infertility. *Urology Research & Practice, 29*, 296–300.

Owaba, A. D., Etim, E. I., Johnson, E. C., & Umoh, U. F. (2021). Aphrodisiac agents used in traditional medicine and their mechanisms of action: A review. *Journal of Pharmacognosy and Phytochemistry, 10*(3), 126–153.

Peterson, B. D., Newton, C. R., & Rosen, K. H. (2016). Examining congruence between partners' perceived infertility-related stress and its relationship to marital adjustment and depression in infertile couples. *Family Process, 55*(2), 239–254.

Popoola, O. O. (2019). Phytochemical status of different solvent extracts of *Anthocleista djalonensis* A. Chev. leaves. *International Annals of Science, 8*(1), 62–69. https://doi.org/10.21467/ias.8.1.62-69

Practice Committee of the American Society for Reproductive Medicine. (2006). Diagnostic evaluation of the infertile male: A committee opinion. *Fertility and Sterility, 86*(5 Suppl. 1), S202–S205.

Practice Committee of the American Society for Reproductive Medicine. (2015). Diagnostic evaluation of the infertile male: A committee opinion. *Fertility and Sterility, 103*(3), e18–e25.

Prakash, D., Singh, B. N., & Mishra, G. (2013). Phytochemicals of *Mucuna pruriens*: A review. *Journal of Medicinal Plants Research, 7*(18), 1285–1298.

Prasad, K. N., Yang, B., & Dong, X. (2006). *Mucuna pruriens*: A comprehensive review. *Pharmacognosy Reviews, 1*(2), 157–162.

Prasad, S., & Aggarwal, B. B. (2011). Turmeric, the golden spice: From traditional medicine to modern medicine. In I. F. F. Benzie & S. Wachtel-Galor (Eds.), *Herbal medicine: Biomolecular and clinical aspects* (2nd ed., Chapter 13). CRC Press.

Puppel, K., Kapusta, A., & Kuczyńska, B. (2015). The etiology of oxidative stress in the various species of animals: A review. *Journal of the Science of Food and Agriculture, 95*(11), 2179–2184. https://doi.org/10.1002/jsfa.7015

Qazi, M., & Molvi, K. (2016). Herbal medicine: A comprehensive review. *International Journal of Pharmaceutical Research, 8*, 1–5.

Ramaiah, M., Chakravathi, G., & Yasaswini, K. (2013). In vitro biological standardization, formulation and evaluation of directly compressed polyherbal anthelmintic tablets. *Pharmacognosy Journal, 5*(3), 130–134.

Ricci, E., Al-Beitawi, S., Cipriani, S., Alteri, A., Chiaffarino, F., Candiani, M., Gerli, S., & Viganò, P. (2020). Semen quality and alcohol intake: A systematic review and meta-analysis. *Reproductive BioMedicine Online, 40*(1), 1–11.

Risberg, K., Fodstad, Ø., & Andersson, Y. (2011). Synergistic anticancer effects of the 2.27PE immunotoxin and ABT-737 in melanoma. *PLOS ONE, 6*(9), e24012. https://doi.org/10.1371/journal.pone.0024012

Rosa, A. C., Corsi, D., Cavi, N., Bruni, N., & Dosio, F. (2021). Superoxide dismutase administration: A review of proposed human uses. *Molecules, 26*(7), 1844. https://doi.org/10.3390/molecules26071844

Sadiku, M. N. O., Chukwu, U. C., Ajayi-Majebi, A., & Musa, S. M. (2022). Traditional medicine in Nigeria. *International Journal of Trend in Scientific Research and Development, 6*(3), 1768–1775.

Savage, G. P. (2010). Velvet bean (*Mucuna pruriens* L.): A versatile crop for the humid tropics. *Tropical and Subtropical Agroecosystems, 12*(1), 107–123.

Sekhar, R. V., Patel, S. G., Guthikonda, A. P., Reid, M., Balasubramanyam, A., Taffet, G. E., & Jahoor, F. (2011). Deficient synthesis of glutathione underlies oxidative stress in aging and can be corrected by dietary cysteine and glycine supplementation. *The American Journal of Clinical Nutrition, 94*(3), 847–853. https://doi.org/10.3945/ajcn.110.003483

Sharma, R. K., Agarwal, A., & Sharma, V. (2013). Assisted reproductive technology and male infertility: Current status and future prospects. *Frontiers in Bioscience—Elite, 5*(2), 897–930.

Sharma, R., Harlev, A., Agarwal, A., & Esteves, S. C. (2016). Cigarette smoking and semen quality: A new meta-analysis examining the effect of the 2010 World Health Organization laboratory methods for the examination of human semen. *European Urology, 70*(4), 635–645. https://doi.org/10.1016/j.eururo.2016.04.010

Shukla, K. K., Mahdi, A. A., Ahmad, M. K., Shankhwar, S. N., Rajender, S., Jaiswar, S. P., & Tiwari, S. C. (2009). *Mucuna pruriens* improves male fertility by its action on the hypothalamus–pituitary–gonadal axis. *Fertility and Sterility, 92*(6), 1934–1940.

Sies, H. (1999). Glutathione and its role in cellular functions. *Free Radical Biology and Medicine, 27*(9–10), 916–921. https://doi.org/10.1016/S0891-5849(99)00177-X

Simmonds, N. W., & Shepherd, K. (1955). The taxonomy and origins of the cultivated bananas. *Journal of the Linnean Society of London, Botany, 55*(359), 302–312. https://doi.org/10.1111/j.1095-8339.1955.tb00015.x

Singh, Z., Karthigesu, I. P., Singh, P., & Kaur, R. (2014). Use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review. *Iranian Journal of Public Health, 43*(Suppl. 3), 7–16.

Suri, S. (2012). Ayurvedic medicinal properties of banana. *CryoLetters, 20*(1), 20–25.

Swennen, R., & Vuylsteke, D. (1993). Banana cultivation in Africa: Diversity, cultivar preferences and cultivation practices. *Outlook on Agriculture, 22*(3), 147–154.

Tainer, J. A., Getzoff, E. D., Richardson, J. S., & Richardson, D. C. (1983). Structure and mechanism of copper, zinc superoxide dismutase. *Nature, 306*(5940), 284–287. https://doi.org/10.1038/306284a0

Tang, S. W., Yao, Y., Andrade, C., & Li, Z. J. (2015). Recent durability studies on concrete structures. *Cement and Concrete Research, 78*, 143–154.

Tavilani, H., Doosti, M., & Saeidi, H. (2005). Malondialdehyde levels in sperm and seminal plasma of asthenozoospermic men and their relationship with semen parameters. *Clinica Chimica Acta, 356*, 199–203.

Thonneau, P., Marchand, S., Tallec, A., Ferial, M. L., Ducot, B., Lansac, J., & Lopes, P. (1991). Incidence and main causes of infertility in a resident population of three French regions, 1988–1989. *Human Reproduction, 6*(6), 811–816.

Tilburt, J. C., & Kaptchuk, T. J. (2008). Herbal medicine research and global health: An ethical analysis. *Bulletin of the World Health Organization, 86*(8), 594–599. https://doi.org/10.2471/BLT.07.042820

Tung, B. T., Nham, D. T., Hai, N. T., & Thu, D. K. (2019). *Curcuma longa*, the polyphenolic curcumin compound and pharmacological effects on liver. In R. R. Watson & V. R. Preedy (Eds.), *Dietary interventions in liver disease* (pp. 125–134). Academic Press.

von Ossowski, I., Hausner, G., & Loewen, P. C. (1993). Molecular evolutionary analysis based on the amino acid sequence of catalase. *Journal of Molecular Evolution, 37*, 71–76.

Wang, C., & Swerdloff, R. S. (2014). Hormonal approaches to male contraception. *Current Opinion in Urology, 24*(6), 514–520.

Wu, G., Fang, Y.-Z., Yang, S., Lupton, J. R., & Turner, N. D. (2004). Glutathione metabolism and its implications for health. *The Journal of Nutrition, 134*(3), 489–492. https://doi.org/10.1093/jn/134.3.489

Yadav, R. P., & Tarun, G. (2017). Versatility of turmeric: A review of the golden spice of life. *Journal of Pharmacognosy and Phytochemistry, 6*(1), 41–46.

Yasui, K., & Baba, A. (2006). Therapeutic potential of superoxide dismutase for resolution of inflammation. *Inflammation Research, 55*, 359–363. https://doi.org/10.1007/s00011-006-5195-y

Youn, H.-D., Kim, E.-J., Roe, J.-H., Hah, Y. C., & Kang, S.-O. (1996). A novel nickel-containing superoxide dismutase from *Streptomyces* spp. *Biochemical Journal, 318*(3), 889–896. https://doi.org/10.1042/bj3180889

Younus, H. (2018). Therapeutic potentials of superoxide dismutase. *International Journal of Health Sciences, 12*(3), 88–93.

Yusuf, A. A., Lawal, B., Alozieuwa, U. B., Onikanni, A. S., Lukman, H. Y., Fadaka, A. O., Olawale, F., Osuji, O., Sani, S., Owolabi, M. S., & Adewuyi, A. H. (2023). Attenuating effects of *Azanza garckeana* fractions on glycemo-impaired-associated dyslipidemia, hepatopathy, and nephropathy. *American Journal of Translational Research, 15*(10), 5997–6014. https://pmc.ncbi.nlm.nih.gov/articles/PMC10641334

Yusuf, A. A., Lawal, B., Sani, S., Garba, R., Mohammed, B. A., Oshevire, D. B., & Adesina, D. A. (2020). Pharmacological activities of *Azanza garckeana* (Goron Tula) grown in Nigeria. *Clinical Phytoscience, 6*, Article 27. https://doi.org/10.1186/s40816-020-00173-0