

Effect of *Jatropha Gossypiifolia* Leaves Extract in Wistar Albino Rats Against Gentamicin-Induced Biochemical Injury

Isaac John Umaru¹, Tensaba Andes Akafa², Joseph Oteng³, Ingwu Joseph Akem⁴, Ocheifa Mathew Ngbede⁵, Ashaka Fidelis Utioukpan⁶, Aboki Nwunuji Mijinyawa⁷, Odok Endurance Akam⁸, Ogholo Ogholo Ekup⁹

^{1,2}Federal University Wukari, Taraba State, Nigeria; ³Ghana Health Service, Public Health Department, Ghana; ⁴Department of Environmental Health Abia State, Nigeria; ⁵Federal Medical Centre, Makurdi, Benue State, Nigeria; ^{6,8}College of Health Technology Calabar, Nigeria; ⁷Taraba State College of Technology Takum, Taraba State, Nigeria; ⁹Open University Nigeria, Calabar, Nigeria
umaru.isaac@fuwukari.edu.ng

Article Info:

Submitted:	Revised:	Accepted:	Published:
Feb 17, 2025	Mar 1, 2025	Mar 13, 2025	Mar 18, 2025

Abstract

Jatropha gossypiifolia L. (Euphorbiaceae), widely known as “bellyache bush,” is a medicinal plant largely used throughout Africa and America. Several human and veterinary uses in traditional medicine are described for different parts and preparations based on this plant. The protective potentials of ethanol leaf extract of *Bryophyllum pinnatum* against gentamicin induced biochemical injury in Wistar albino rats were evaluated using serum biochemical parameters. Twenty-five male albino rats were divided randomly into five groups A to E. Groups C and D received 100, 200 and 300 mg/kg *Jatropha gossypiifolia* leaves extract respectively. Groups A and B were fed with chow (normal and positive controls) while Group E received 100 mg/kg of Vitamin C. Biochemical injury was induced in rats in groups B-E with gentamicin (i.p) at the dose of 80 mg/kg body weight on the 16th day of study. The rats were

then fasted for 48 h and sacrificed by cervical dislocation. Serum was collected for biochemical analysis using standard methods and analytical biochemical kits. There was a significant increase in the serum levels of AST, ALT and ALP of the rats administered with gentamicin (Group B - positive control) compared to the Group A- normal control. However, treatment of rats with 100, 200 and 300 mg/kg body weight of ethanol leaf extract of *Jatropha gossypifolia* L significantly decrease these biochemical parameters compared to Group B ($p < 0.05$). Also, there was a significant increase in the serum levels of conjugated bilirubin and total bilirubin in Group B compared to the normal control. Rat pre-treated with 200 and 300 mg/kg body weight of extract showed significant decrease in the conjugated and total bilirubin compared to the non-treated rats in Group B ($p < 0.05$). Serum level of urea and creatinine significantly increased in Group B compared to normal control. Rats in groups C and D pre-treated with 200 and 300 mg/kg body weight of extract showed significant decrease in the levels of urea and creatinine compared to Group B ($p < 0.05$). There was significant decrease in the serum levels of albumin and total protein of the rats administered with gentamicin compared to the normal control. Rats pre-treated with 200 and 300 mg/kg body weight of extract showed significant increase compared rats in non-treated Group B ($p < 0.05$). The results obtained from this study shows that the ethanolic leaf extract of *Jatropha gossypifolia* L has protective functions against gentamicin-induced hepatic and nephrotic damage in Wistar albino rats.

Keywords: Effect, *Jatropha Gossypifolia*, Leaves, Wistar Albino, Rats, Gentamicin, Induced Biochemical, Injury

INTRODUCTION

The Euphorbiaceae family, which is considered one of the largest families of Angiosperms, covers about 7,800 species distributed in approximately 300 genera and 5 subfamilies worldwide. These species occur preferentially in tropical and subtropical environments [1, 2].

Among the main genera belonging to this family, there is *Jatropha* L., (figure 1) which belongs to the subfamily Crotonoideae, Jatropeae tribe and is represented by about 200 species. This genus is widely distributed in tropical and subtropical regions of Africa and the Americas [1]. The name “*Jatropha*” is derived from the Greek words “*jatros*,” which means “doctor” and “*trophe*,” meaning “food,” which is associated with its medicinal uses [3]. The *Jatropha* genus is divided into two subgenera, *Jatropha* and *curcas*, from which the subgenus *Jatropha* has the widest distribution, with species found in Africa, India, South

America, West Indies, Central America, and the Caribbean [4]. *Jatropha* species are used in traditional medicine to cure various ailments in Africa, Asia, and Latin America or as ornamental plants and energy crops [3]. Several known species from genus *Jatropha* have been reported for their medicinal uses, chemical constituents, and biological activities such as *Jatropha curcas*, *Jatropha elliptica*, *Jatropha gossypifolia*, and *Jatropha mollissima*, among others [3].

From these species, *Jatropha gossypifolia* L. (Figure 1) is discussed here. It is a vegetal species widely known as “bellyache bush” and is a multipurpose medicinal plant largely used in folk medicine for the treatment of various diseases [3, 5, 6]. It is widely distributed in countries of tropical, subtropical, and dry tropical weather and tropical semiarid regions of Africa and the Americas [7].

In Brazil, it predominates in the Amazon, Caatinga, and Atlantic Forest and is distributed throughout the country in the North, Northeast, Midwest, South, and Southeast regions [8]. Several human and veterinary uses in traditional medicine are described for different parts (leaves, stems, roots, seeds, and latex) and preparations (infusion, decoction, and maceration, among others) based on this plant, by different routes (oral or topical). The most frequent reports concern its antihypertensive, anti-inflammatory, antiophidian, analgesic, antipyretic, antimicrobial, healing, antianemic, antidiabetic, and antihemorrhagic activities, among many other examples [3, 5, 7, 9]. Other uses are also related to this plant, such as biodiesel production, pesticide, insecticide, vermifuge, ornamentation, and even its use in religious rituals [3, 6, 10–13].

An important feature of *J. gossypifolia* species is that, due to its important potential medicinal applications, it is included in the National List of Medicinal Plants of Interest, which includes 71 species of medicinal plants that have the potential to generate pharmaceutical products of interest to public health [14].

Regarding its phytochemical constitution, alkaloids, coumarins, flavonoids, lignoids, phenols, saponins, steroids, tannins, and terpenoids were already detected in different extracts from different parts of this plant [15]. Among the main activities already studied for this species (including various types of extracts from different parts of the plant), the antihypertensive, antimicrobial, anti-inflammatory, antioxidant, and antineoplastic activities mainly stand out, supporting some of its popular uses [3, 16].

Some toxicity studies have shown that despite the known toxicity of *Jatropha* species, *J. gossypifolia* presented low toxicity in some in vitro and in vivo experiments. However, some studies have indicated that ethanolic extract from the leaves, in acute oral use, is safe for rats, but with chronic use, it could be toxic [17–19].

The leaf juice of the plant was used traditionally as antiviral, antipyretic, antimicrobial, anti-inflammatory, antitumor, hypocholesterolemic, antioxidant, diuretic, antiulcer, antidiabetic, antiseptic, cough suppressant, antihistamine and anti-allergic (Okwu and Josiah, 2006).

The plant has also been used for the treatment of oedema of legs [20]. Leaf juice is used in the treatment of coughs, bronchial affections, blood dysentery, jaundice and gout (Ghani, 2003). In South-eastern Nigeria, the herb is used to facilitate the expulsion of the placenta after delivery and applied on the body of young children when they are sick [21].

Gentamicin is an aminoglycoside antibiotic widely used for the treatment of bacterial infections. Therapeutic doses of gentamicin and other aminoglycoside antibiotics can produce nephrotoxicity in humans and animals. Its use is known as one of the most common causes of acute renal failure and nephrotoxicity [22]. Vitamin C, selenium, vitamin E, taurine and the carotenoids (betacarotene, lutein and lycopene) were reported to decrease the gentamicin-induced reduction in the glomerular filtration rate and the severity of the tubular damage [23,24].

So, in view of the potential applications of this plant, this study aims to provide an up-to-date on the effect of *Jatropha gossypifolia* leaves in wistar albino rats against gentamicin-induced biochemical injury.



Figure 1: *Jatropha gossypifolia* plant showing the leaves and the fruits

Table 1: Phytochemical Composition of *Jatropha Gossypifolia*; Methanol, Aqueous Extract, Stem Bark, Roots-Bark and Leaves

S/N	Phytochemicals	<i>Jatropha gossypifolia</i>				
		Methanol	Aqueous	Stem bark	Roots bark	Leaves
1	Tannins	+	+	+	-	+
2	Alkaloids	+	+	-	-	+
3	Phenols	+	+	+	+	+
4	Flavonoids	+	+	+	+	+
5	Terpenoids	-	+	+	+	+
6	Glycosides	+	+	+	+	-
7	Antraquinones	+	+	+	+	-
8	Saponin	+	+	+	+	+
9	Cyanoglycoside	+	+	+	+	-
10	Cardiac glycosides	+	+	+	+	+
11	Phlobatannin	+	+	+	+	+

MATERIALS AND METHODS

Animals: Twenty-five Wistar Albino male rats weighing 110 to 1320g were obtained from the Animal House of the HAUEMM Animal Farm, Federal Housing Estate Bajabure Gerie. Adamawa State. The animals were housed in stainless cages and maintained under standard conditions for the period of the study. The animals were acclimatized for two weeks and maintained at the temperature of $25 \pm 2^\circ\text{C}$, $45 \pm 5\%$ relative humidity and 12 h light/dark cycle. The animals had access to water and food freely.

Chemicals

Absolute methanol was product of BDH Chemical Company 164 Ltd, Poole, England. Rat chow was purchased from Pfizer Nigeria Plc.

Collection of Plant Materials

The plant *Jatropha Gossypifolia* was collected from HAUEMM Animal Farm, Federal Housing Estate Bajabure Gerie. Adamawa State

Extract Preparation

The leaves of *Jatropha Gossypifolia* were collected and air dried under shade for two weeks. The dried leaves were ground into powder using an electric blender. 600 g of the powdered leaf were macerated in 1800 ml of ethanol at room temperature for 24 h. It was continuously mixed and then filtered using a filter paper (Whatman size No.1). The filtrate was dried in a water bath at 37°C , and the 50 g viscous concentrate was kept in air tight bottle at 4°C until use.

Experimental Design

Twenty-five (25) adult male Wistar albino rats (eight weeks old) were randomly divided into five (5) groups of five (5) animals per group, labelled A to E and were treated as follows:

1. Group A: served as the normal control (fed and water only).
2. Group B: served as the positive control.
3. Group C: received 100 mg/kg body weight of extract orally.
4. Group D: received 200 mg/kg body weight of extract orally.
5. Group E: received 300 mg/kg body weight Vitamin C orally (Standard antioxidant).

All treatments lasted for fifteen (15) days. Biochemical injury in animals in Group B to E were induced with gentamicin (i.p) at the dose of 80 mg/kg body weight on the 16th day of study, fasted for 48 h and then sacrificed through cervical dislocation.

Collection of Samples

Blood samples were collected in plain bottles through the cardiac puncture and were centrifuged at 2,800 rpm for 10 min. Serum obtained was used for biochemical analysis.

Biochemical Assay

Serum transaminase (ALT and AST) was determined by method of Reitman and Frankel (1957). ALP by the phenolphthalein monophosphate method (Babson, 1965). Total protein was determined by colorimetric method (Biuret method), as modified by Gornaliet et al. (1994) method. Bilirubin was estimated by colorimetric method of Jendrassik and Grof (1938). Serum Urea was estimated by Natelson (1951) method, and serum creatinine by Jelliffe (1971).

Statistical Analysis

The results were expressed as mean \pm SD. Data was analyzed by one-way analysis of variance (ANOVA). Sequential differences among means were calculated at the level of $P < 0.05$, using Turkey contrast analysis as needed.

RESULTS

Table 2. Effect of *Jatropha Gossypifolia* on Liver Indices in Gentamicin-Induced Wistar Albino Rats.

Group	JATROPHA GOSSYPIIFOLIA ON LIVER INDICES							
	Treatment (Extract)	AST (IU/L)	ALT (IU/L)	ALP (IU/L)	Albumin (g/dl)	Total Protein (g/dl)	Conj. Bilirubin (μ mol/l)	Total Bilirubin (μ mol/l)
A	Normal Control	20.23 \pm 2.94a	18.00 \pm 1.58a	40.19 \pm 2.10a	36.89 \pm 1.00c	64.36 \pm 1.36c	1.55 \pm 0.17a	5.76 \pm 0.64a
B	Positive Control	26.61 \pm 6.06c	30.20 \pm 4.76c	44.19 \pm 1.55a	24.19 \pm 1.55a	45.73 \pm 2.65a	1.84 \pm 0.15b	9.22 \pm 1.42b
C	100mg/kg bwt	23.22 \pm 2.38b	22.40 \pm 4.04b	44.24 \pm 5.43b	34.24 \pm 5.43b	58.97 \pm 3.71b	1.60 \pm 0.23a	6.37 \pm 1.24a
D	200mg/kg bwt	22.63 \pm 2.19b	23.00 \pm 4.30b	43.46 \pm 6.23b	36.89 \pm 1.00c	57.42 \pm 4.72b	1.56 \pm 0.13a	6.39 \pm 0.73a
E	300mg/kg bwt	22.23 \pm 2.38b	22.80 \pm 4.15b	40.15 \pm 6.50b	24.19 \pm 1.55a	55.83 \pm 1.83b	1.47 \pm 0.16a	6.25 \pm 0.42a

Data are mean \pm SD (n = 5). Means values in the same column with different superscript letter(abc) are significantly different; $p < 0.05$

Table 3. Effect of *Jatropha Gossypifolia* on Kidney Indices in Gentamicin-Induced Wistar Albino Rats.

Group	JATROPHA GOSSYPIIFOLIA ON KIDNEY INDICES		
	Treatment (Extract)	UREA (mmol/l)	CREATININE (mmol/l)
A	Normal Control	5.34 ± 0.51a	69.41 ± 3.47a
B	Positive Control	9.13 ± 0.37b	92.37 ± 6.71c
C	100mg/kg bwt	6.29 ± 0.92a	76.27 ± 7.46b
D	200mg/kg bwt	6.11 ± 1.12a	76.35 ± 6.12b
E	300mg/kg bwt	6.36 ± 0.41a	73.27 ± 5.57a

Data are mean ± SD (n = 5). Means values in the same column with different superscript letter(s) are significantly different; p < 0.05

The results of the effect of ethanol leaf extract of *Jatropha Gossypifolia* on liver indices in wistar rats are presented in Table 1. There was significant increase in serum levels of AST, ALT, ALP of the rats administered with gentamicin (26.61 ± 6.06c, 30.20 ± 4.76c, 24.19 ± 1.55a), respectively when compared to the normal control group (20.23 ± 2.94a, 18.00 ± 1.58a, 40.19 ± 2.10a), (p < 0.05). However, pre-treatment of rats with 200 and 300 mg/kg body weight of Methanol leaf extract of *Jatropha Gossypifolia* significantly decrease AST, ALT, ALP (22.63 ± 2.19b, 23.00 ± 4.30b, 30.46 ± 6.23b) and (22.23 ± 2.38b, 22.80 ± 4.15b, 33.15 ± 6.50b), respectively, compared to non-treated positive control group (p < 0.05).

However, there was a significant increase in the serum levels of conjugated bilirubin and total bilirubin in non-treated rat positive control group (1.84 ± 0.15b and 9.22 ± 1.42b), respectively, compared to the normal control (1.55 ± 0.17a, and 5.76 ± 0.64a). Rats in the groups treated with 200 and 300 mg/kg body weight of extract showed significant decrease in the conjugated and total bilirubin (1.56 ± 0.13a and 6.39 ± 0.73a) and (1.47 ± 0.16a, and 6.25 ± 0.42a), respectively, compared to the non-treated positive control group (p < 0.05).

The serum level of urea and creatinine significantly increased in gentamicin induced rats' group (9.13 ± 0.37b and 92.37 ± 6.71c), respectively, compared to normal control (5.34 ± 0.51a and 69.41 ± 3.47a), (p < 0.05) (Table 2). Rat in groups D and E treated with 200 and 300 mg/kg body of extract showed significant decrease in the levels of urea and creatinine (6.11 ± 1.12a and 76.35 ± 6.12b) and (6.36 ± 0.41a and 73.27 ± 5.57a), respectively, compared to the positive control group (p < 0.05).

There was a significant decrease in the serum levels of albumin and total protein of the rats administered with gentamicin ($24.19 \pm 1.55a$ and $45.73 \pm 2.65a$), respectively, compared to the normal control ($36.89 \pm 1.00c$, and $64.36 \pm 1.36c$).

Rats treated with 200 and 300 mg/kg body weight of extract showed significant increase ($36.89 \pm 1.00c$ and $57.42 \pm 4.72b$) and ($24.19 \pm 1.55a$ and $55.83 \pm 1.83b$), respectively, compared to rats in non-treated group ($p < 0.05$).

DISCUSSION

Many researchers have directed their efforts towards the provision of empirical proof to back up the use of many tropical plants for trado-medical practices [26,27, 28]. This study was undertaken to explore the hepatoprotective and nephroprotective effect of *Jatropha Gossypiifolia* leaf extract in the hepatic and nephrotic damage caused by the administration of gentamicin. The plant was observed to have a potential with strong protein oxidation inhibitory potency. This could be attributed to the antioxidant activity of extracts from *J. gossypiifolia* which was evaluated by Kharat et al. [29].

In this work the high content of phenols, tannins, and flavonoids in the leaves prompted the authors to evaluate the antioxidant activity of the leaves. DPPH free radical, ferric thiocyanate, and nitric oxide scavenging methods were used to analyze the antioxidant activity in vitro of methanol, ethyl acetate, and aqueous extracts, demonstrating positive results. The authors attributed the free radical scavenging activity to the presence of flavonoids [30]. On the other hand, a study showed that different extracts (petrol ether, chloroform, ethyl acetate, and n-butanol) from whole plant of *J. gossypiifolia* had only partial antioxidant activity in DPPH scavenging, total antioxidant capacity, and lipid peroxidation tests. Among them, the Methanol and ethyl acetate extract was the most active, which correlates positively with its higher content of phenolic compounds in comparison with the other extracts [30]. Thus, the decreased levels of serum albumin and total proteins in non-treated rat group, is an indication of hepatotoxicity [31].

Albumin is essential for tissue growth and aids in preventing the leakage of fluids from blood vessels. It plays an important role in transporting both endogenous and exogenous substances, serving as protein reserves, as well as maintaining osmotic pressure [32]. However, there was significant increase in the levels of albumin and total protein in extract treated rat groups.

The present study corroborates suggestions made by Davies and Goldberg, [33] that the herb *Jatropha Gossypifolia* leaf extract may have strong protein oxidation inhibitory potency and thus, may be a good source of medicines against diseases in which lipids and protein oxidation are involved, such as toxic hepatitis since protein degradation seems to occur by distinct mechanism [34]. The significant increase in serum level of urea and creatinine in gentamicin induced rats' group is also an indication of nephrotoxicity.

Therapeutic doses of gentamicin and other aminoglycoside antibiotics can produce nephrotoxicity in humans and animals and use of this class of antibiotics is known as one of the most common causes of acute renal failure possibly due to increased renal uptake of the antibiotic mainly by the proximal tubules. Nephrotoxicity is the most common side effects associated with the use of gentamicin [35]. Rats in groups treated with extract showed significant decrease in the levels of urea and creatinine. Kanika, [36] and Nwali et al. [37], reported some phytochemical constituents of *Jatropha Gossypifolia* leaf extract to include; vitamin E, selenium, vitamin C, taurine and the carotenoids (beta-carotene, lutein and lycopene) which have the potentials to decrease the gentamicin-induced reduction in the glomerular filtration rate and the severity of the tubular damage [38, 39].

The aqueous extract of *Jatropha Gossypifolia* leaves was earlier reported to possess potent nephroprotective activity in gentamicin induced nephrotoxicity in rats. The results obtained from this study show that the methanolic leaf extract of *Jatropha Gossypifolia* leaf extract has protective function against gentamicin-induced Albino rats.

CONCLUSION

As demonstrated by this study, *J. gossypifolia* presents an important potential for the generation of pharmacological and/or biotechnological products, based on popular uses and biological studies scientifically showing its properties. However, regarding specifically its medicinal properties, further studies are still necessary to assay important folk uses of the species and characterize the major compounds responsible for the bioactivity. Thus, the results obtained from this study show that the ethanolic leaf extract of *Jatropha Gossypifolia* leaves has protective function against gentamicin-induced damage in rats.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

REFERENCES

1. G. L. Webster, "Classification of the euphorbiaceae," *Annals of the Missouri Botanical Garden*, vol. 81, pp. 3–143, 1994.
2. M. V. Alves, "Checklist das espécies de Euphorbiaceae Juss. ocorrentes no semi-árido pernambucano, Brasil," *Acta Botânica Brasileira*, vol. 12, no. 3, pp. 485–495, 1998.
3. C.W. Sabandar, N. Ahmat, F. M. Jaafar, and I. Sahidin, "Medicinal property, phytochemistry and pharmacology of several *Jatropha* species (Euphorbiaceae): a review," *Phytochemistry*, vol. 85, pp. 7–29, 2013.
4. C. K. A. Leal and M. D. F. Agra, "Estudo farmacobotânico comparativo das folhas de *Jatropha molissima* (Pohl) Baill," *Acta Farmaceutica Bonaerense*, vol. 24, no. 1, pp. 5–13, 2005.
5. L. C. Di Stasi and C. A. Hiruma-Lima, *Plantas medicinais na Amazônia e na Mata Atlântica*, UNESP, São Paulo, Brazil, 2nd edition, 2002.
6. M. D. F. Agra, K. N. Silva, I. J. L. D. Basílio, P. F. De Freitas, and J.M. Barbosa-Filho, "Survey of medicinal plants used in the region Northeast of Brazil," *Brazilian Journal of Pharmacognosy*, vol. 18, no. 3, pp. 472–508, 2008.
7. S. R. Mariz, A. C. R. Borges, M. F. F. Melo-Diniz, and I. A. Medeiros, "Possibilidades terapêuticas e riscos toxicológicos de *Jatropha gossypifolia* L.: uma revisão narrativa," *Revista Brasileira De Plantas Medicinai*s, vol. 12, no. 3, pp. 346–357, 2010.
8. I. Cordeiro and R. Secco, "*Jatropha gossypifolia* L. Lista de espécies da flora do Brasil," *Jardim Botânico do Rio de Janeiro*, <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB17581>.
9. U. P. de Albuquerque, P.M. de Medeiros, A. L. S. de Almeida et al., "Medicinal plants of the caatinga (semi-arid) vegetation of NE Brazil: a quantitative approach," *Journal of Ethnopharmacology*, vol. 114, no. 3, pp. 325–354, 2007.
10. U. P. Albuquerque, L. H. C. Andrade, and J. Caballero, "Structure and floristics of homegardens in Northeastern Brazil," *Journal of Arid Environments*, vol. 62, no. 3, pp. 491–506, 2005.
11. S. A. Ceasar and S. Ignacimuthu, "Applications of biotechnology and biochemical engineering for the improvement of *Jatropha* and Biodiesel: a review," *Renewable and Sustainable Energy Reviews*, vol. 15, no. 9, pp. 5176–5185, 2011.
12. U. P. de Albuquerque, J.M. Monteiro, M. A. Ramos, and E. L. C. de Amorim, "Medicinal and magic plants from a public market in northeastern Brazil," *Journal of Ethnopharmacology*, vol. 110, no. 1, pp. 76–91, 2007.
13. A. Falodun, Q. Sheng-Xiang, G. Parkinson, and S. Gibbons, "Isolation and characterization of a new anticancer diterpenoid from *Jatropha gossypifolia*," *Pharmaceutical Chemistry Journal*, vol. 45, no. 10, pp. 636–639, 2012.
14. BRASIL, "Fitoterapia: plantas de interesse ao SUS," Brazilian Health Ministry, <http://portal.saude.gov.br/portal/arquivos/pdf/RENISUS.pdf>.

15. X.-P. Zhang, M.-L. Zhang, X.-H. Su, C.-H. Huo, Y.-C. Gu, and Q.-W. Shi, "Chemical constituents of the plants from genus *Jatropha*," *Chemistry and Biodiversity*, vol. 6, no. 12, pp. 2166–2183, 2009.
16. S. K. Sharma and H. Singh, "A review on pharmacological significance of genus *Jatropha* (Euphorbiaceae)," *Chinese Journal of Integrative Medicine*, vol. 18, no. 11, pp. 868–880, 2012.
17. S.R.Mariz, G. S.Cerqueira,W. C. Ara'ujo et al., "Chronic toxicologic study of the ethanolic extract of the aerial parts of *Jatropha gossypifolia* in rats," *Revista Brasileira de Farmacognosia*, vol. 22, no. 3, pp. 663–668, 2012.
18. S. R.Mariz, M. S. T. Araujo, G. S. Cerqueira et al., "Histopathological evaluation in rats after acute treatment with the ethanol extract from aerial parts of *Jatropha gossypifolia* L.," *Revista Brasileira De Farmacognosia*, vol. 18, no. 2, pp. 213–216, 2008.
19. S. R. Mariz, G. S. Cerqueira, W. C. Ara'ujo et al., "Estudo toxicol'ogico agudo do extrato etan'olico de partes a'ereas de *Jatropha gossypifolia* L. em ratos," *Revista Brasileira de Farmacognosia*, vol. 16, no. 3, pp. 372–378, 2006.
20. Okwu DE, Josiah C (2006). Evaluation of the chemical composition of two Nigerian medicinal plants, *Afr. J. Biotechnol.*, 5(4): 357-361.
21. Okwu DE, Nnamdi FU (2011). Two novel flavonoids from *Bryophyllum pinnatum* and their antimicrobial Activity. *Pharm. Chem. J.*, 3(2):1-10.
22. Agoha RC (1974). Medicinal Plants of Nigeria, Offset Drakkerij Faculfcitder Wiskunde in Naturwetenschappen,the Netherlands 33:41.
23. Cuzzocrea S, Mazzon E, Dugo L, Serraino L, Paola RD, Britti D, Sarro Pierpaoli S, Caputi AP, Masini E, Savemini D (2002). A role of superoxide in gentamicin mediated nephropathy in rats. *Eur. J. Pharmacol.*, 54(5): 469-473
24. Anganeyulu M, Chopra K (2004). Effect of irbesartan on the antioxidant defense system and nitric oxide release diabetic rat kidney. *Am. J. Nephrol.*, 24(5): 488-496.
25. Ekor M, Farombi EO, Emerole GO (2006). Modulation of gentamicin induced renal dysfunction and injury by the phenolic extract of soybean (*Glycine max*). *Fundam. Clin. Pharmacol.*, 20(3): 263-271.
26. Ojo, O. A., Ajiboye, B. O., Ojo, A. B., Oyinloye, B. E., Imiere, O. D., & Adeyonu, O. (2017). Ameliorative potential of *Blighia sapida* KD Koenig bark against pancreatic β -cell dysfunction in alloxan-induced diabetic rats. *Journal of complementary and Integrative medicine*, 14(3), 20160145.
27. Sule, O. J., & Arhoghro, M. E. (2016). Protective potentials of *Bryophyllum pinnatum* in Wistar albino rats against gentamicin-induced biochemical injury. *Journal of Medical and Biological Science Research*, 2(10), 163-167.
28. Madusolomuo MA, Okoye ZS (1995). Anti-coagulant properties of bergenin from *sacoglottis gabonsis* stem bark extract *J. Med Red.*, 23:443-444.
29. [55]. A. R. Kharat, A. K. Dolui, and S. Das, "Free radical scavenging potential of *Jatropha gossypifolia*," *Asian Journal of Chemistry*, vol. 23, no. 2, pp. 799–801, 2011.
30. [48]. D. Shahwar, S.-U. Shafiq, N. Ahmad, S. Ullah, and M. A. Raza, "Antioxidant activities of the selected plants from the family Euphorbiaceae, Lauraceae, Malvaceae and Balsaminaceae," *African Journal of Biotechnology*, vol. 9, no. 7, pp. 1086–1096, 2010.
31. Abatan MO, Arowolo ROA, Olorunsogo O (1996). Pathological Effects of *Lantana camara* and *Dichapetalum madagascasiense* in Goats. *Trop. Vet. Med.*,14:127 -132.

32. Kpomah ED, Arhoghro EM, Uwakwe AA (2012). Biochemical and histopathological changes following chronic administration of diherbal mixture of zanthoxylum lepreurii and piper guineense J. Nat. Sci. Res. 2(6):22-28
33. Davies KJA, Goldberg AL (1987). Oxygen radicals stimulate intracellular proteolysis and lipid peroxidation by independent mechanisms in erythrocytes. J. Biol. Chem., 262(17): 8220-8226.
34. Njyou N, Fredric Paul F, Moundipa Angele N, Tchana Bonaventure, Ngadjui T Felicite M(2008). Tchouanguiep Inhibition of microsomal lipid peroxidation and protein oxidation by extracts from plants used in BAMUN folk medicine (Cameroon) against hepatitis. Afr. J. Trad., Complem. Alternative Med., 5(3):278-289.
35. Cuzzocrea S, Mazzon E, Dugo L, Serraino L, Paola RD, Britti D, Sarro Pierpaoli S, Caputi AP, Masini E, Savemini D (2002). A role of superoxide in gentamicin mediated nephropathy in rats. Eur. J. Pharmacol., 54(5): 469-473
36. Kanika P (2011). Pharmacognostic & Phytochemical Evaluation of Bryophyllum pinnatum Leaves. J. Adv. Sci. Res., 2(1): 42-49.
37. Nwali BU, Okaka ANC, Ibiam UA, Aja PM (2012). Phytochemical composition of Bryophyllum pinnatum Leaves. Int. J. Adv. Biol. Res., 2(4): 614-616.
38. Anganeyulu M, Chopra K (2004). Effect of irbesartan on the antioxidant defense system and nitric oxide release diabetic rat kidney. Am. J. Nephrol., 24(5): 488-496.
39. Ekor M, Farombi EO, Emerole GO (2006). Modulation of gentamicin induced renal dysfunction and injury by the phenolic extract of soybean (Glycine max). Fundam. Clin. Pharmacol., 20(3): 263-271.