

### Major Depressive Disorder (MDD): A Multifactorial Mental Health Crisis and Approaches to Holistic Management

Muhammad Akram<sup>1</sup>, Isaac John Umaru<sup>2</sup>, Abid Mahmood<sup>3</sup>, Mohammed Khudhair Hasan<sup>4</sup>, Hind A. Abdulghafoor<sup>5</sup>, Fahad Said Khan<sup>6</sup>, Fethi Ahmet Ozdemir<sup>7</sup>, Gaweł Sołowski<sup>8</sup>, Jaouher Ben Ali<sup>9</sup>, Jehan Mohammed Al-Musawi<sup>10</sup>

<sup>1,3</sup>Government College University Faisalabad, Pakistan; <sup>2</sup>Federal University Wukari, Taraba State, Nigeria; <sup>4</sup>Al Manara College for Medical Sciences, Maysan, Iraq; <sup>5</sup>University of Fallujah, Iraq; <sup>6</sup>University of Poonch Rawalakot, Azad Jammu and Kashmir, Pakistan;

<sup>7,8</sup>Bingol University, Bingol, 1200, Türkiye; <sup>9</sup>University of Tunis, Tunisia;

<sup>10</sup>University of Kufa, Najaf, Iraq

makram\_0451@yahoo.com

#### Article Info:

Submitted:	Revised:	Accepted:	Published:
Apr 7, 2025	Apr 21, 2025	May 3, 2025	May 8, 2025

#### Abstract

MDD, also known as Major Depressive Disorder, or clinical depression, is one of the most common and debilitating mental illnesses worldwide since it affects a person's emotional, cognitive, and physiological functioning balance. Sufferers typically experience unrelenting low mood, anhedonia or lack of interest in activities, extreme tiredness, and numerous loopy mental and physical symptoms that negatively affect normal functioning. It's worth emphasizing that a complex disorder, not solely an emotional state, depression arises as a result of the interplay of genetics, predisposition to the illness, neurobiological underpinnings, psychosocial and environmental stressors, as well as their myriad factors. This paper analyzes the causes, symptoms, diagnostic guidelines, and the most recent methods of treatment available for MDD. Special focus is given to psychotherapy, pharmacologic approaches, lifestyle changes, and the necessity for a holistic approach that integrates all dimensions of patient care.

**Keywords:** Major Depressive Disorder, depression, serotonin, mental health, CBT, antidepressants, neurobiology, mood disorders, suicide prevention, psychological therapy

## Introduction

In today's world, one of the most important public health issues is depression. The WHO says that more than 280 million people in the world suffer from Major Depressive Disorder (MDD), which indicates a serious break in emotional and functional capabilities which is far more intense than mere sadness or grieving (Strakowski et al 2015). Unlike temporary melancholic spells, MDD is enduring and can precipitate severe deterioration in social, professional, academic and personal activities. Its impact is far-reaching as it increases the burden of disability and unemployment, substance dependency, and suicidal tendencies. Depression is not associated with age, gender, or background. Even though it is so widespread, distressing and misrepresentation is still prevalent which unacceptably delays detection and treatment. The burden on people and society can be lowered significantly with accepting depression as a medical issue.

## Etiology and Risk Factors

The development of MDD arises from multiple factors which include biological, psychological, sociocultural, and genetic factors (Remes et al 2021). Neurobiological theories indicate that depression results from underactivity of certain neurotransmitters, especially serotonin and dopamine, and norepinephrine, which are important for mood, motivation, and energy. Some people with MDD show structural and functional changes in the prefrontal cortex, amygdala, and hippocampus which indicate poor regulation of emotions, along with impaired cognitive processes. Depressive disorder has a genetic component; increased risk is observed in first degree relatives, although other factors are nonspecific. Social and psychological aspects are particularly important; prolonged abuse, trauma, neglect during early life or significant loss produces predisposition towards depression later on. Other known triggers include chronic stress, low social support, and poor socioeconomic status. From the cognitive point of view, a person suffering from depression tends to have negative self evaluation along with pessimistic view of the future which Beck described as cognitive triad (Beck et al 2016). Factors that heighten the risk of

developing MDD also include hormonal changes, chronic illness, like thyroid and cardiovascular disorders, and substance abuse.

### **Symptoms and Diagnostic Criteria**

A Major Depressive Disorder (MDD) is marked by a low persistent mood or an absence of interest or pleasure in almost all activities for a minimum of two weeks (Giannelli et al 2020). Based on DSM-5, depression will be diagnosed if at least five of the following symptoms are present along with one core symptom (depressed mood or anhedonia):

1. Fatigue or loss of energy
2. Feelings of worthlessness or excessive guilt
3. Impaired concentration or indecisiveness
4. Insomnia or hypersomnia
5. Psychomotor agitation or retardation
6. Significant weight loss or gain
7. Recurrent thoughts of death or suicide

The symptoms must not be attributable to a medical condition, substance use, or other reason and must cause significant distress or impairment in functioning (Schuckit et al 2006). There is a wide range in the severity and course of depression. A person may experience a single episode or face chronic and recurrent forms. Important subtypes with distinct clinical characteristics encompass atypical depression, melancholic features, seasonal patterns, or postpartum depression.

### **Impact on Functioning and Society**

The impacts of depression extend well beyond the individual's personal pain and suffering; it harms overall productivity and economic factors like healthcare spending (Donohue et al 2007). In depressed employees, productivity is affected because of absenteeism, presenteeism (working while unwell), and reduced performance. Adolescents with MDD show lack of motivation, poor attention, and difficulty with social interactions, diminishing their academic performance as well. On the sociocultural spectrum, depression increases vulnerability to becoming homeless, incarcerated and to socially withdrawing from society.

Perhaps most tragically, suicide, which ranks as the leading cause of death claiming 700,000 lives each year, is fundamentally linked with untreated depression. Mental illness remains one of the most stigmatized issues today, making it even harder for people to seek the help they need (Corrigan et al 2014). Understanding depression as a medical problem rather than a flaw or weakness enables addressing the issue at both individual and system levels.

### **Treatment Approaches**

To successfully resolve Major Depressive Disorder (MDD), both medical and non-medical treatments need to work in synergy. The initial approach to managing depression is through medication (Dowrick et al 2009). The two most prescribed types of MDD medications are SSRIs (selective serotonin reuptake inhibitors) and SNRIs (serotonin-norepinephrine reuptake inhibitors). Examples include fluoxetine and/or sertraline for SSRIs and venlafaxine and/or duloxetine for SNRIs. These medications increase the level of key neurotransmitters linked to mood in the brain, therefore enhancing the overall mental state of the patient. Psychotherapy, particularly CBT (cognitive behavioral therapy) and IPT (interpersonal therapy), is often combined with medication as it has proven highly effective for moderate to severe depression cases. In CBT, patients are taught to recognize and change harmful patterns of thinking and behaving. IPT emphasizes restoring healthy social functioning in patients and improving their interpersonal relationships (Lipsitz et al 2013). Additionally, mindfulness-based cognitive therapy (MBCT) helps prevent recurring relapses in patients with a history of multiple depressive episodes. In certain instances where diagnosis does not respond to treatment, electro-convulsive therapy (ECT), Transcranial Magnetic Stimulation (TMS), or fiberglass ketamine infusion therapy have offered greater success. Alongside professional help, adopting healthy lifestyle changes that include exercising, well-balanced diets, sufficient rest, and minimal drug and alcohol consumption greatly aid in the recovery process. Moreover, online platforms offering mental health services have expanded the range of available public resources.

### **Prevention and Long-Term Management**

Even though not all types of occurring depression could be managed or mitigated, early intervention and oversight can severely minimize its severity and frequency. Identifying depression in greater risk groups such as adolescents, new mothers, and chronic illness

patients can enable prompt diagnosis and treatment (Sawyer et al 2007). Effective public educational campaigns positively influence help-seeking behavior and change the prevailing attitude on mental health. These campaigns have proven to significantly benefit the society. Sustaining recovery from depression calls for comprehensive support, guidance, and treatment of other concurrent illnesses which may include anxiety, substance use disorder, and personality disorders. The holistic approach allows the patient to formulate active coping mechanisms. The use of strong social networks and emotional resilience also aid the patient (Musich et al 2019). It is undeniably important to highlight that recovery programs that help in maintaining decreed medication along with the adapting to changes in lifestyle, ongoing therapy, as well as those aimed at stabilizing remission, can greatly enhance life quality.

## **Conclusion**

Major depressive disorders (MDD) emerge as one of the most common mental disorders affecting several people at the same time. Common depression occurs due to biological, psychological and social change and results in low self-esteem and social avoidance. Advanced techniques help us understand the importance of treating depression through therapies that engage medication and psychosocial support. Critical attention such conditions face helps in treating it with utmost care in a society with stigma. Every individual suffering from depression or any mental ailment needs timely diagnosis and the right treatment plan to avoid social isolation. Change in mindset enables better focus on risk education and international movement towards treatment.

## **References**

- Beck, A. T., & Bredemeier, K. (2016). A Unified Model of Depression: Integrating Clinical, Cognitive, Biological, and Evolutionary Perspectives. *Clinical Psychological Science*, 4(4), 596–619.
- Corrigan, P. W., Druss, B. G., & Perlick, D. A. (2014). The Impact of Mental Illness Stigma on Seeking and Participating in Mental Health Care. *Psychological Science in the Public Interest*, 15(2), 37–70.
- Donohue, J. M., & Pincus, H. A. (2007). Reducing the Societal Burden of Depression: A Review of Economic Costs, Quality of Care and Effects of Treatment. *Pharmacoeconomics*, 25, 7–24.

Muhammad Akram, Isaac John Umaru, Abid Mahmood, Mohammed Khudhair Hasan, Hind A. Abdulghafoor, Fahad Said Khan, Fethi Ahmet Ozdemir, Gawel Solowski, Jaouher Ben Ali, Jehan Mohammed Al-Musawi

- Dowrick, C. (2009). *Beyond Depression: A New Approach to Understanding and Management*. Oxford University Press.
- Giannelli, F. R. (2020). Major Depressive Disorder. *JAAPA*, 33(4), 19–20.
- Lipsitz, J. D., & Markowitz, J. C. (2013). Mechanisms of Change in Interpersonal Therapy (IPT). *Clinical Psychology Review*, 33(8), 1134–1147.
- Musich, S., Wang, S. S., Slindee, L., Kraemer, S., & Yeh, C. S. (2019). Association of Resilience and Social Networks with Pain Outcomes Among Older Adults. *Population Health Management*, 22(6), 511–521.
- Remes, O., Mendes, J. F., & Templeton, P. (2021). Biological, Psychological, and Social Determinants of Depression: A Review of Recent Literature. *Brain Sciences*, 11(12), 1633.
- Sawyer, S. M., Drew, S., Yeo, M. S., & Britto, M. T. (2007). Adolescents with a Chronic Condition: Challenges Living, Challenges Treating. *The Lancet*, 369(9571), 1481–1489.
- Schuckit, M. A. (2006). Comorbidity Between Substance Use Disorders and Psychiatric Conditions. *Addiction*, 101, 76–88.
- Strakowski, S., & Nelson, E. (2015). *Major Depressive Disorder*. Oxford University Press.