

## **Application of Cauchy's Residue Theorem in Evaluating Real and Trigonometric Integrals**

**Ndam G. S., Manjak N. H., Adamu M. S., Kpop S. S.**

Abubakar Tafawa Balewa University, Bauchi, Nigeria

ndamgeorgeshehu@gmail.com

### **Article Info:**

| <b>Submitted:</b> | <b>Revised:</b> | <b>Accepted:</b> | <b>Published:</b> |
|-------------------|-----------------|------------------|-------------------|
| Jun 18, 2026      | Jul 16, 2026    | Jul 28, 2026     | Aug 2, 2026       |

---

### **Abstract**

Evaluating definite real and trigonometric integrals remains a fundamental problem in mathematical analysis, particularly when elementary calculus techniques are insufficient or inefficient. This study aims to examine the application of Cauchy's Residue Theorem as an analytical method for evaluating selected definite real and trigonometric integrals. The analysis transforms the original integrals into appropriate contour integrals in the complex plane, identifies the relevant isolated singularities, computes their residues, and applies the Residue Theorem to derive exact integral values. The results show that this approach provides systematic and exact solutions to integrals that are otherwise difficult to evaluate using conventional real-variable methods. The study concludes that residue theory offers an effective and mathematically elegant framework for definite integral evaluation. These findings reinforce the practical significance of complex analysis and provide a methodological reference for selecting suitable contours and calculating residues in advanced mathematical problem-solving.

**Keywords:** Complex Analysis; Contour Integration; Definite Integrals; Residue Theorem; Trigonometric Integrals

## Introduction

Complex analysis is a branch of mathematics that studies functions of a complex variable. One of its most important tools is Cauchy's Residue Theorem, which provides a powerful technique for evaluating contour integrals. Residue theory has numerous applications in mathematics, physics, and engineering.

Many definite real and trigonometric integrals that are difficult to solve using elementary methods can be evaluated efficiently using residue calculus. The purpose of this paper is to demonstrate the application of Cauchy's Residue Theorem in solving selected real and trigonometric integrals.

## Preliminary Concepts

### Isolated Singularities

A point  $z_0$  is called an isolated singularity of a function  $f(z)$ , if  $f(z)$  is analytic in a neighborhood of  $z_0$  except at  $z_0$  itself.

### Residue

The residue of a function  $f(z)$  at an isolated singularity  $z_0$  is the coefficient of  $(z - z_0)^{-1}$  in its Laurent series expansion.

### Cauchy's Residue Theorem

Theorem:

Let  $f(z)$  be analytic inside and on a simple closed contour  $c$  except for finitely many isolated singularities  $z_1, z_2, \dots, z_n$ , inside  $c$ .

Then

$$\int_c f(z) dz = 2\pi i \sum \text{Res}(f, z_k).$$

Where  $\text{Res}(f, z_k)$  denotes the residue of  $f(z)$  at  $z_k$ .

### Application of Cauchy's Residue Theorem to Real and Trigonometric Integrals

This section demonstrates the application of Cauchy's Residue Theorem in evaluating selected real and trigonometric integrals. The procedure consists of transforming each integral into a contour integral, identifying the singularities enclosed by the chosen contour, calculating the corresponding residues, and applying the Residue Theorem to obtain the desired result.

Note: Let  $c$  denote the unit circle  $|z| = 1$ .

#### Evaluate

$$I = \int_{-\infty}^{\infty} \frac{dx}{x^2 + 1}.$$

#### Solution

Define the complex function

$$f(z) = \frac{1}{z^2 + 1}.$$

The simple pole occur at

$$z = \pm i.$$

Choose a semicircular contour  $c_R$  in the upper half-plane consisting of the interval  $[-R, R]$  and the semicircular arc of radius  $R$ .

The only pole enclosed by  $c_R$  is

$$z = i.$$

Since the pole is simple,

$$\text{Res}(f, i) = \lim_{z \rightarrow i} (z - i) \frac{1}{(z - i)(z + i)} = \frac{1}{2i}.$$

By Cauchy's Residue Theorem,

$$\int_{C_R} f(z) dz = 2\pi i \left( \frac{1}{2i} \right) = \pi.$$

As  $R \rightarrow \infty$ , the contribution from the semicircular arc tends to zero by Jordan's Lemma.

Therefore,

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} = \pi.$$

Hence,

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} = \pi.$$

**Evaluate**

$$I = \int_{-\infty}^{\infty} \frac{\cos(ax)}{x^2 + 1} dx, \quad a > 0.$$

**Solution**

By cosine function

$$\cos(ax) = \operatorname{Re}(e^{iax}).$$

Thus consider

$$f(z) = \frac{e^{iaz}}{z^2 + 1}.$$

The poles occur at

$$z = \pm i.$$

Only  $z = i$ , lies in the upper half-plane.

The residue at  $z = i$  is

$$\operatorname{Res}(f, i) = \lim_{z \rightarrow i} (z - i) \frac{e^{iaz}}{(z - i)(z + i)} = \frac{e^{-a}}{2i}.$$

Applying the Residue Theorem,

$$\int_{C_R} \frac{e^{aiz}}{z^2 + 1} dz = 2\pi i \left( \frac{e^{-a}}{2i} \right) = \pi e^{-a}.$$

Taking the real part yields

$$\int_{-\infty}^{\infty} \frac{\cos(ax)}{x^2 + 1} dx = \pi e^{-a}, \quad \forall a > 0.$$

**Evaluate**

$$I = \int_0^{2\pi} \frac{d\theta}{5 + 4 \cos \theta}.$$

**Solution**

Let  $z = e^{i\theta}.$

Then  $d\theta = \frac{dz}{iz},$

And  $\cos \theta = \frac{z + z^{-1}}{2}.$

By substituting into the integral gives

$$I = \int_c \frac{dz}{5 + 4\left(\frac{z + z^{-1}}{2}\right)iz}.$$

We have

$$I = \int_c \frac{dz}{i(2z^2 + 5z + 2)}.$$

The poles are to be obtain by factoring

$$2z^2 + 5z + 2 = 0.$$

$$(2z + 1)(z + 2) = 0.$$

Hence,

$$z = -\frac{1}{2}, \quad z = 2.$$

Only

$$z = -\frac{1}{2},$$

lies inside the unit circle.

Thus the residue is

$$\operatorname{Res}\left(\frac{1}{i(2z+1)(z+2)}, -\frac{1}{2}\right) = \frac{1}{3i}.$$

By applying the Residue theorem,

$$I = 2\pi i \left(\frac{1}{3i}\right) = \frac{2\pi}{3}.$$

Therefore,

$$\int_0^{2\pi} \frac{d\theta}{5 + 4\cos\theta} = \frac{2\pi}{3}.$$

**Evaluate**

$$I = \int_0^{\infty} \frac{dx}{x^4 + 1}.$$

**Solution**

The poles of

$$f(z) = \frac{1}{x^4 + 1}$$

are given as

$$z^4 = -1.$$

Thus

$$z = e^{\frac{i\pi}{4}}, \quad e^{\frac{3i\pi}{4}}, \quad e^{\frac{5i\pi}{4}}, \quad e^{\frac{7i\pi}{4}}.$$

The poles in the upper half-plane are

$$z = e^{\frac{i\pi}{4}}, \quad \text{and} \quad e^{\frac{3i\pi}{4}}.$$

By computing the residues and applying the Residue Theorem gives

$$\int_{-\infty}^{\infty} \frac{dx}{x^4 + 1} = \frac{\pi}{\sqrt{2}}.$$

Since the integrand is even as,

$$\int_0^{\infty} \frac{dx}{x^4 + 1} = \frac{\pi}{2\sqrt{2}}.$$

## Discussion of Results

The examples demonstrate that Cauchy's Residue Theorem provides a systematic method for evaluating both real and trigonometric integrals. The principal advantage of the method lies in reducing integral evaluation to residue computation at isolated singularities. In each case, the choice of contour and identification of poles lead directly to exact closed form results. These examples highlight the efficiency and elegance of complex variable techniques compared with traditional real variable techniques.

## Conclusion

This study has demonstrated the usefulness of Cauchy's Residue Theorem in solving real and trigonometric integrals. The theorem provides a direct and elegant method for evaluating integrals through contour integration and residue computation. The examples considered illustrate the effectiveness of complex analysis in solving problems that arise in mathematical analysis and related fields.

## References

- Ahlfors, L. V. (1979). *Complex analysis: An introduction to the theory of analytic functions of one complex variable* (3rd ed.). McGraw-Hill.
- Brown, J. W., & Churchill, R. V. (2014). *Complex variables and applications* (9th ed.). McGraw-Hill Education.
- Conway, J. B. (1978). *Functions of one complex variable I* (2nd ed.). Springer-Verlag. <https://doi.org/10.1007/978-1-4612-6313-5>
- Gamelin, T. W. (2001). *Complex analysis*. Springer. <https://doi.org/10.1007/978-0-387-21607-2>
- Marsden, J. E., & Hoffman, M. J. (1999). *Basic complex analysis* (3rd ed.). W. H. Freeman.
- Needham, T. (2023). *Visual complex analysis* (25th anniversary ed.). Oxford University Press. <https://global.oup.com/academic/product/visual-complex-analysis-9780192868923>
- Rudin, W. (1987). *Real and complex analysis* (3rd ed.). McGraw-Hill.
- Spiegel, M. R., Lipschutz, S., Schiller, J. J., & Spellman, D. (2009). *Schaum's outline of complex variables* (2nd ed.). McGraw-Hill. <https://www.mheducation.com/highered/mhp/product/schaum-s-outline-complex-variables-2ed.html>