

## GC-MS PROFILING AND PATHOGENIC EFFECT OF KHAYA ANTHOTHECA METHANOLIC LEAVES EXTRACT ON SOME SELECTED FUNGI

Isaac John Umaru<sup>1</sup>, Philip Shadrach<sup>2</sup>, Godwin Baaku Adi<sup>3</sup>, Egeonu Stephen  
Ugoeze<sup>4</sup>, Chakfa Nanmar<sup>5</sup>, Akafa Andes Tansaba<sup>6</sup>, Kerenhappuch Isaac Umaru<sup>7</sup>,  
and Bando Christopher David<sup>8</sup>

<sup>1,2,3,4,5,6</sup>Federal University Wukari, Taraba State, Nigeria

<sup>7</sup>University of Higher Institute Buea, South West Cameroon, Cameroon

<sup>8</sup>National Biotechnology Research and Development Agency, Jalingo, Nigeria  
umaruisaac@gmail.com

### Article Info:

|              |             |              |              |
|--------------|-------------|--------------|--------------|
| Submitted:   | Revised:    | Accepted:    | Published:   |
| May 20, 2024 | Jun 7, 2024 | Jun 10, 2024 | Jun 14, 2024 |

### Abstract

This research evaluated GC-MS profile, and pathogenic effect of methanolic leaves extracts of Khaya anthotheca on Some Selected fungi. Khaya anthotheca a member of Meliaceae family is traditionally used for treating several ailments. The leaves were harvested in Sukundi road Wukari and air dried for four weeks, the leaves were pulverized into powder using manual blender and stored in an air tight container. Extraction was done with absolute methanol; the extract was filtered using mesh and the filtrate was concentrated at 68°C. Only two grams of the filtrate were used for the GC-MS study. One-way analysis of variance (ANOVA) was used for statistical analysis. The GC-MS result revealed 50 constituents. Result of Methanolic extracts of leaves of

Khaya anthotheca only showed significant activity against *Aspergillus niger* at dose 100 µg/mL, *Candida tropicalis* and *Fusarium oxysporium* at dose 500 µg/mL. This study supports the use of Khaya anthotheca by indigenous herbalists by providing scientific proof of the plant's ability to treat pathogenic diseases. To establish the effectiveness of this plant in the management of fungal infections, more clinical trials at the clinical levels are needed.

**Keywords:** Antifungal, Khaya Anthotheca, Phytochemicals, Medicinal

## INTRODUCTION

The rising prevalence of pathogenic microorganisms which are resistant to the modern antimicrobial drugs that have been produced in the last three decades [1, 4, 5] This has been a threat to health as a result of the high cost, adulteration and increasing toxic side effects of these synthetic drugs coupled with their inadequacy in diseases treatment found more especially in the developing countries cannot be over emphasized [20]. Among the pathogenic microorganism fungal diseases is one of the most threatening and major cause of morbidity and mortality worldwide. [2] As a result of the number of multi-drug resistant microbial strains and the appearance of strains with reduced susceptibility to antibiotics continuously increasing. This increase has been attributed to indiscriminate use of broad-spectrum antifungal and immunosuppressive agents [3, 4]. Coincidentally, the last decade has also witnessed increasing intensive studies on extracts and biologically active compounds isolated from plant species used for natural therapies or herbal medicine. [5, 7]

Medicinal plants represent a rich source of antimicrobial agents. Plants are used medicinally in different countries and are a source of many potent and powerful drugs [8]. A wide range of medicinal plant parts is used for extract as raw drugs and they possess varied medicinal properties. While some of these raw drugs are collected in small quantities by the local communities and traditional healers for local use [19]. Considering the vast potentiality of plant as sources for antimicrobial drugs with reference to antifungal agents, an investigation as regard the plant Khaya anthotheca was undertaken to screen the local flora for its antifungal activity. This situation provided the impetus to the search for new antifungal substances from various sources like Khaya anthotheca. Traditional medicine

has made use of many different plant extracts for treatment of fungal infections and some of these have been tested for in vitro antifungal activity [6].



**Figure 1:** *Khaya anthotheca* leaf

*Khaya anthotheca* is a member of the family Meliaceae. Six species make up this little genus; two are found in Madagascar and the Comores and four are found in tropical Africa [8]. The paripinnate leaves and mostly spherical, 4-5 valved, dehiscent woody capsules make the genus easily identifiable [8]. In addition to being frequently used for window frames, paneling, doors and stair cases, the wood is highly prized for use in furniture, cabinetry, veneer and decorative boxes and cases. It works well for light flooring, automotive bodies, musical instruments, sporting goods, toys, plywood, pulpwood, shipbuilding and carving [9].

Research also demonstrated that *Khaya anthotheca* possesses neuroprotective action [10]. Some important bioactive constituents such as gedunin and limonoids which are of great benefits have been reported to be present in this plant [11]. Yacine et al.[12] also reported the presence of phytochemicals such as sterols and polyterpenes, flavonoids, polyphenols, saponins and tannins in the plant. Elegant research also reported that *K. anthotheca* possesses high antioxidant and antiradical properties due to the presence of bioactive

constituents such as polyphenols, polyterpenes and in addition to tannins and saponins<sup>13</sup>. Studies also documented that *K. anthotheca* possesses antimalarial [7], anti-protozoal [14], anticancer [15] and neuroprotective [16] activities. However, reports on the anti-ulcerogenic and anti-bacterial efficacy of plants are rather rare or speculative. This study therefore evaluates the anti-ulcerogenic and anti-bacterial activities of methanolic crude extract of *Khaya anthotheca* root on indomethacin treated albino rats

## METHODS

### Materials

Mortar and pestle, beakers, Conical flask, Spatula, Filter paper (Whatman No.1), Masking tape, Hand gloves, Cotton wool, face mask, Separating funnel, Digital analytical weighing balance (OHAUS: PA-4120), Refrigerator (HTF-146F), Thermostatic Water Cabinet (MODEL: HH-W420), Micro pipette, Test tubes and Test tube racks, and sample bottles, Glucometer (Accu chek)

### Chemicals and Reagents

Methanol, distilled water, 75% of alcohol, were purchased from Justlab Nigeria Limited 140 Isolo road Egbe - Okotun, Lagos - Nigeria. All chemicals used were of highest quality.

### Plant collection and identification

The *Khaya anthotheca* Leaves was rinsed under running water and air dried. To prevent ultraviolet light from eliminating the active ingredients, the leaves were air-dried under shade [18,19]. Using a mortar and pestle, the dried leaves were ground into a fine powder and kept in an airtight container until required.

### Determination of % yield of extract

The percentage yield of the extract of *Khaya anthotheca* was determined by weighing the coarse sample before extraction and the *Khaya anthotheca* ethanol leaves extract after concentration and then calculated using the formula.

### GC- MS phytochemical profiling

The GC-MS analysis was performed in a 7890A gas chromatograph that also functions as a mass spectrometer (Agilent 19091-433HP, USA). It is equipped with a triple axis detector and an inert MSD 5675c infection as well as an HP-5 MS fused silica column (5% phenyl

methyl siloxane, 30.0m 250m). Helium gas was set to flow at a rate of 1.0 ml per minute as a carrier gas. Other GC-MS conditions include a 1  $\mu$ L injector in split mode with a split ratio of 1:50 and an injection temperature of upto 300 °C. Ion source temperature is 250 °C, interface temperature is 300 °C, pressure is 16.2 psi, and out time is 1.8 mm. Over the course of five minutes, the column's temperature rose from 36°C to 150°C at a rate of 4°Cmin<sup>-1</sup>. At a rate of 20°Cmin<sup>-1</sup>, the temperature increased to 250°C and maintained there for 5 minutes. Elution lasted for 47.5 minutes in total. By comparing each component's average peak area to the sum of all areas, the corresponding percentage quantity of each component was computed. The supplier's MS solution software was utilized to manage the system and gather data. Mass Hunter Library NIST17 was used in comparing and identifying the constituents.

### **Antifungal potential**

The antifungal potential of the plant extract was performed by agar disc diffusion method as reported by [20, 21].

### **Fungal preparation**

The fungi were standardized by extracting fungal spores from a 7-day-old culture on SDA slant and introducing a normal saline solution that was sterile with a 48-hour pure culture. Turbidity was adjusted to match 0.5 McFarland stand standards. With the help of sterilized glass beads, ten milliliters of pure normal saline containing 3% w/v Tween 80 was employed to distribute the spores. A UV spectrophotometer (Spectronic 20D; Milton Roy Company, Pacisa, Madrid, Spain) set to a transmittance of 70–72% of the suspensions was used to standardize the spore suspension to 1.0106 spores/mL. The prepared plates were allowed to incubate for 24 hours at 37°C [22].

## RESULTS AND DISCUSSION

### Phytochemical constituents of leaves of *Khaya anthotheca* methanol crude extract


**Table 1:** Phytochemical profile of *leaves Khaya anthotheca* methanol crude extract

| S/N | Name                                          | R.Time | Area      | Height   |
|-----|-----------------------------------------------|--------|-----------|----------|
| 1   | Dodecane                                      | 15.326 | 3444740   | 1271238  |
| 2   | Tetradecane                                   | 18.304 | 150967299 | 16523371 |
| 3   | Undecanal                                     | 18.600 | 2087082   | 894718   |
| 4   | 1-Undecanol                                   | 20.355 | 2595232   | 841001   |
| 5   | Tetradecane                                   | 20.848 | 4828945   | 1819690  |
| 6   | Dodecanal                                     | 21.378 | 77701897  | 15172477 |
| 7   | (E)-.beta.-Famesene                           | 22.304 | 1629324   | 613674   |
| 8   | 1-Dodecanol                                   | 23.007 | 46136232  | 10172333 |
| 9   | Benzene, 1-(1,5-dimethyl-4-hexenyl)-4-meth    | 23.182 | 8868440   | 3176963  |
| 10  | Heneicosane                                   | 23.444 | 51540066  | 13142990 |
| 11  | .alpha.-Farnesene                             | 23.643 | 2003129   | 791455   |
| 12  | .beta.-Bisabolene                             | 23.825 | 2834162   | 1117573  |
| 13  | Tridecanal                                    | 23.936 | 61459079  | 13774706 |
| 14  | (1S,5S)-4-Methylene-1-((R)-6-methylhept-5-    | 24.263 | 1923373   | 697160   |
| 15  | 1,6,10-Dodecatrien-3-ol, 3,7,11-trimethyl-, ( | 25.098 | 5000845   | 1357406  |
| 16  | 1-Tetradecanol                                | 25.594 | 58505167  | 9246835  |
| 17  | Heneicosane                                   | 25.815 | 2017498   | 509524   |
| 18  | Caryophyllene oxide                           | 25.923 | 7120718   | 1860590  |
| 19  | Tetradecanoic acid, 2-oxo-, methyl ester      | 26.215 | 1502359   | 334904   |
| 20  | Tetradecanal                                  | 26.678 | 416219832 | 22244814 |
| 21  | Neointermedeol                                | 26.986 | 12224325  | 1199765  |
| 22  | cis-1-Chloro-9-octadecene                     | 27.861 | 114555022 | 13733228 |
| 23  | Heneicosane                                   | 28.140 | 42557088  | 6691262  |
| 24  | Pentadecanal-                                 | 28.665 | 19732438  | 6276275  |
| 25  | Dodecanal, O-methyloxime                      | 28.790 | 2878675   | 171426   |
| 26  | 1-Tetradecanol                                | 28.992 | 6256826   | 864056   |
| 27  | 2-((2R,4aR,8aS)-4a-Methyl-8-methylenedeca     | 29.291 | 5442959   | 890324   |
| 28  | 9,12-Octadecadienoic acid (Z,Z)-              | 29.440 | 1449467   | 347279   |
| 29  | Tetradecanoic acid                            | 29.955 | 78259929  | 7209241  |
| 30  | Tetradecanoic acid, 2-oxo-, methyl ester      | 30.225 | 9026018   | 1022046  |
| 31  | cis-9-Hexadecenal                             | 30.415 | 4910683   | 727269   |
| 32  | Eicosanal-                                    | 30.874 | 45458691  | 10063730 |
| 33  | 2-Pentadecanone, 6,10,14-trimethyl-           | 31.277 | 31303258  | 4888448  |
| 34  | 2-Methyltetracosane                           | 31.590 | 3024460   | 351320   |
| 35  | Pentadecanoic acid                            | 31.862 | 15252037  | 2175323  |
| 36  | n-Pentadecanol                                | 32.171 | 8432271   | 2324597  |
| 37  | Heneicosane                                   | 32.371 | 30492415  | 5915702  |
| 38  | 5,9,13-Pentadecatrien-2-one, 6,10,14-trimeth  | 32.762 | 7336599   | 1090178  |
| 39  | Hexadecanoic acid, methyl ester               | 32.969 | 6012999   | 1408015  |
| 40  | Andrographolide                               | 33.183 | 8706779   | 2327089  |
| 41  | Bicyclo[4.4.0]dec-6-en-9.beta.-ol, 1,7-dimeth | 33.363 | 12182572  | 1734591  |



|    |                                               |        |           |          |
|----|-----------------------------------------------|--------|-----------|----------|
| 42 | n-Hexadecanoic acid                           | 34.428 | 376367103 | 11986924 |
| 43 | Docosanoic acid, 1,2,3-propanetriyl ester     | 34.865 | 27302036  | 1338321  |
| 44 | Bicyclo[2.2.1]heptan-2-ol, 3-amino-1,7,7-tri  | 35.296 | 9873042   | 1131538  |
| 45 | Oleic Acid                                    | 35.691 | 15580728  | 2525909  |
| 46 | n-Hexadecanoic acid                           | 35.982 | 24635421  | 3842420  |
| 47 | 7-Hexadecenal, (Z)-                           | 36.372 | 27120743  | 5193798  |
| 48 | 8,11,14-Docosatrienoic acid, methyl ester     | 36.516 | 7910817   | 2105619  |
| 49 | Phytol                                        | 36.804 | 68746295  | 10546910 |
| 50 | 9-Octadecenoic acid, 1,2,3-propanetriyl ester | 38.163 | 406333014 | 10069983 |

The GC-MS results of methanolic extracts of *leaves Khaya anthotheca* revealed tetradecanal, tetradecane, 1-undecanol, pentadecanoic acid, neointermedeol, 1-dodecanol, heneicosane, docosanoic acid 1,2,3-propanetriyl ester, phytol, 9-octadecenoic acid, 1,2,3-propane triyl ester, 1-tetradecanol and andrographolide etc. as some of the constituents.



**Figure 2:** GC-MS chromatogram of *Khaya anthotheca* methanol leaves crude extract

Result obtained from GC-MS of methanolic extracts of *leaves of Khaya anthotheca* in table 1. Fifty (50) constituents were identified. Tetradecane is a long-chain fatty aldehyde in which two hydrogen atoms connected to a terminal carbon are changed by an oxo group to form tetradecanal. It serves as a metabolite for both plants and microorganisms. It is both a 2,3-saturated and long-chain fatty aldehyde. It comes from a tetradecane hydride [23]. A common monounsaturated fat in the human diet is oleic acid. Oral consumption of monounsaturated fat is associated with lower levels of LDL cholesterol and maybe higher

levels of HDL cholesterol [24,25]. Oleic acid is a crucial omega-9 monounsaturated fatty acid that is used as an emulsifier. It has also been reported to be hypotensive [26] and to slow the advancement of adrenoleukodystrophy, a dangerous condition that affects the brain and adrenal glands [27,28]. The manufacturing of personal care and cosmetic products uses n-hexadecenoic acid, which has also been shown to have antioxidant, hypocholesterolemia, anti-inflammatory, and 5-alpha reductase inhibitor effects [29–30]. Antioxidant, neuroprotective, antibacterial, anticancer, anti-inflammatory and anti-diuretic properties of phytol have been documented [28,31]. The anti-inflammatory properties of pentadecanoic acid are excellent [32].

1-undecanol increases male attractiveness in sex pheromone trap species *Grapholita molesta* (Busck) [33]. 1-Dodecanol is a fatty alcohol that is dodecane with a hydroxy group in place of hydrogen from one of the methyl groups. It is approved for use as Lepidopteran pheromone/sex bait in apple and pear orchards to prevent particular insects, the larvae of which destroy crops, from mating. It serves as a pheromone, pesticide, plant metabolite, insect attractant and cosmetic [23]. Heneicosane is used as an antimicrobial [34]. Tetradecanoic acid has been demonstrated to exhibit antifungal action against the fungi *C. albicans* and *A. niger* when combined with hydrazide-hydrazones [35]. Administration of *Khaya anthotheca* leaves extract to animals or humans may exert some the pharmacological activities discussed on the active phytochemicals present in this extract.

Antifungal activity of *Khaya anthotheca*

**Table 2:** Antifungal activity of *Khaya anthotheca* in on *Aspergillus niger*, *Aspergillus flavus*, *Candida tropicalis* and *Fusarium oxysporium* in millimetre (mm)

| Solvent | Organism                  | controls        |             | Extracts millimeter (mm) |                 |                |                |
|---------|---------------------------|-----------------|-------------|--------------------------|-----------------|----------------|----------------|
|         |                           | Nystatin+v<br>e | DMSO<br>-ve | 50                       | 100             | 250            | 500            |
| Methan  | <i>Aspergillus niger</i>  | 24.67±0.11      | 30 +<br>0.8 | 12.43±0.1<br>2           | 13.62±0.25<br>* | 12.49±0.1<br>8 | 11.43±<br>0.25 |
|         | <i>Aspergillus flavus</i> | 23.40±0.05      | 30 +<br>0.8 | 10.40±0.1<br>3           | 11.578±0.1<br>8 | 12.57±0.1<br>3 | 12.63±<br>0.15 |



|    |                            |            |             |                |            |                |                 |
|----|----------------------------|------------|-------------|----------------|------------|----------------|-----------------|
| ol | <i>Candida tropicalis</i>  | 23.10±0.08 | 30 +<br>0.8 | 12.17±0.2<br>3 | 12.72±0.12 | 15.47±0.3<br>6 | 16.55±0.11<br>* |
|    | <i>Fusarium oxysporium</i> | 23.20±0.10 | 30 +<br>0.8 | 12.45±0.2<br>6 | 13.20±0.11 | 14.29±0.1<br>8 | 17.60±0.13<br>* |

Result are expressed as mean  $\pm$  standard deviation (N = 5)

\*= significant activity is seen when compared to the control ( $p < 0.05$ )

Concentration of standard is 500  $\mu\text{g/mL}$  of nystatin, Conc= Concentration, DCM = Dichloromethane, DMSO = Dimethyl sulfoxide.

Result of antifungal activity of methanolic extracts of leaves *Khaya anthotheca* showed that application of all doses of extract showed no significant activity ( $p > 0.05$ ) against *Aspergillus niger* except 100  $\mu\text{g/mL}$  which showed significant activity ( $p < 0.05$ ) against *Aspergillus niger* compares to the normal control. Application of all doses of methanolic extracts of leaves *Khaya anthotheca* showed no significant activity against *Aspergillus flavus* when compared to the control. Application of all doses of methanolic extracts of leaves *Khaya anthotheca* showed no significant activity against ( $p > 0.05$ ) *Candida tropicalis* except 500  $\mu\text{g/mL}$  which showed significant activity ( $p < 0.05$ ) against *Candida tropicalis*. Application of all doses of methanolic extracts of leaves *Khaya anthotheca* showed no significant activity ( $p > 0.05$ ) against *Fusarium oxysporium* except 500  $\mu\text{g/mL}$  which showed significant activity ( $p < 0.05$ ) against *Fusarium oxysporium*. The antifungal activity exhibited by *Khaya anthotheca* leaves could due to the presence of phytoconstituent such as phytol, oleic acid, octadecanoic acid etc. The result of antifungal activity suggests that methanolic extracts of leaves *Khaya anthotheca* may have the potential to be used as an antifungal drug [36].

## CONCLUSION

According to the findings in this study, the *Khaya anthotheca* leaves fraction has antifungal activities with less negative effects when administered at low concentrations. *Khaya anthotheca* leaf fraction may be used as a potential effective agent as antifungal which is less expensive, easily accessible, and site-harming because of its antifungal activity displayed by methanolic extract of *Khaya anthotheca* leaves. The outcome of this study show that the plants investigated possess antifungal activities, thus justifying their use in folk medicine for

the treatment of skin and other related infections. long-term treatment with *Khaya anthotheca* methanolic leaves extract may achieve long-term potentials of the extract.

### **Ethics approval and consent to participate**

The study was approved by the Animal Experimentation Ethics Committee of Federal University Wukari, Nigeria. Reference Number AEEC/FUW/014/2023

### **Acknowledgement**

Not applicable.

### **Funding**

This research received no specific grant from any funding agency.

### **Conflict of Interest**

All authors declare that they have no conflict of interest

### **Abbreviations**

RBG, random blood glucose; Conc, concentration; DCM, dichloromethane; DMSO, dimethyl sulfoxide; GCMS, gas chromatography mass spectrophotometer, STZ, streptozotocin; Bwt; body weight.

## **REFERENCES**

1. Akinnuga AM, Bamidele O, Ebunlomo OA, Adeniyi OS, Adeleye GS, Ebomuche LC. (2010). Hypoglycaemic effects of dietary intake of ripe and unripe lycopersicon esculentum (tomatoes) on streptozotocin-induced diabetes mellitus in rats. *Online J Biol Sci.* 10:50-3.
2. El-Serag HB, Tran T, Everhart JE. (2010). Diabetes increases the risk of chronic liver disease and hepatocellular carcinoma. *Gastroenterology.* 2004 Feb 1;126(2):460-8. Available at: <https://doi.org/10.1053/j.gastro.2003.10.065>
3. De Marco R, Locatelli F, Zoppini G, Verlato G, Bonora E, Muggeo M. (1999). Cause-specific mortality in type 2 diabetes. The Verona Diabetes Study. *Diabetes care.* May 1;22(5):756-61. Available at: <https://doi.org/10.2337/diacare.22.5.756>
4. Umar A, Ahmed QU, Muhammad BY, Dogarai BB, Soad SZ. (2010). Anti-hyperglycemic activity of the leaves of *Tetracera scandens* Linn. Merr. (Dilleniaceae) in alloxan induced diabetic rats. *Journal of Ethnopharmacology.* Aug 19;131(1):140-5. Available at: <https://doi.org/10.1016/j.jep.2010.06.016>
5. Brownlee M. (2005). The pathobiology of diabetic complications: a unifying mechanism. *diabetes.* Jun 1;54(6):1615-25. Available at <https://doi.org/10.2337/diabetes.54.6.1615>

6. Albano E. (2006). Alcohol, oxidative stress and free radical damage. *Proceedings of the nutrition society.* Aug;65(3):278-90. Available at: <https://doi.org/10.1079/PNS2006496>
7. Novo E, Parola M. (2008). Redox mechanisms in hepatic chronic wound healing and fibrogenesis. *Fibrogenesis & tissue repair.* Dec;1(1):1-58. Available at: <https://doi.org/10.1186/1755-1536-1-5>
8. Rendell M. (2004). The role of sulphonylureas in the management of type 2 diabetes mellitus. *Drugs.* Jun;64:1339-58. Available at: <https://doi.org/10.2165/00003495-200464120-00006>
9. Das K, Tiwari RK, Shrivastava DK. (2010). Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *Journal of medicinal plants research.* Jan 18;4(2):104-11. Available at: <https://doi.org/10.5897/JMPR09.030>
10. Umaru IJ, Badruddin FA, Umaru HA. (2018). Phytochemical, antifungal and antibacterial potential of *Leptadenia hastata* stem-bark extract. *MOJ Toxicol.*4(4):263-8.
11. Aboh MI, Olayinka BO, Adeshina GO, Oladosu P. (2014). Antifungal activities of phyto compounds from *Mitracarpus villosus* (Sw.) DC from Abuja, Nigeria. *J Microbiol Res.*4(2):86-91. Available at: <https://doi.org/10.5923/j.microbiology.20140402.07>
12. Umaru I, Fasihuddin A, Zaini B. (2017). Antifungal activity of *Leptadenia hastata* (Pers) Decne leaves extract. *Int. J. Pure Appl. Biosci.* 5:14-8. Available at: <http://dx.doi.org/10.18782/2320-7051.5996>
13. National Center for Biotechnology Information. (2023). PubChem Compound Summary for CID 31284, Methyl Tetradecanoate. Retrieved May 10. Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/Methyl-tetradecanoate>.
14. Rao BG, Rao ES, Rao TM. (2012). Quantification of phytochemical constituents and in-vitro antioxidant activity of *Mesua ferrea* leaves. *Asian Pacific Journal of Tropical Biomedicine.* Feb 1;2(2):S539-42. Available at: [https://doi.org/10.1016/S2221-1691\(12\)60269-X](https://doi.org/10.1016/S2221-1691(12)60269-X)
15. Imo C, Yakubu OE, Imo NG, Udegbumam IS, Onukwugha OJ. (2018). Chemical composition of *Xylopia aethiopica* fruits. *Am J Physiol Biochem Pharmacol.*7(2):48-53. Available at: <https://doi.org/10.5455/ajpbp.20180521064020>
16. Mahato SB, Sen S. (1997). Advances in triterpenoid research, 1990–1994. *Phytochemistry.* Apr 1;44(7):1185-236. Available at: [https://doi.org/10.1016/S0031-9422\(96\)00639-5](https://doi.org/10.1016/S0031-9422(96)00639-5)
17. Nestel P, Clifton P, Noakes M. (1994). Effects of increasing dietary palmitoleic acid compared with palmitic and oleic acids on plasma lipids of hypercholesterolemic men. *Journal of lipid research.* Apr 1;35(4):656-62. Available at: [https://doi.org/10.1016/S0022-2275\(20\)41179-4](https://doi.org/10.1016/S0022-2275(20)41179-4)
18. Shadrach P, Imo C, Otitoju O, Mohammad ZI, Asaph PM. (2023). Phytochemical Constituents of Ethanolic Extracts of Riped and Unripened Fruits of *Dennettia tripetala*. *Interna J of Pharma*02.5(1): 01-07.
19. Aparna V, Dileep KV, Mandal PK, Karthe P, Sadasivan C, Haridas M. (2012). Anti-inflammatory property of n-hexadecanoic acid: structural evidence and kinetic

- assessment. *Chemical biology & drug design*. Sep;80(3):434-9. Available at: <https://doi.org/10.1111/j.1747-0285.2012.01418.x>
20. Bruce SO, Onyemalu VO, Orji CE. (2021). Evaluation of the antiulcer activity and GC-MS spectroscopic Analysis of the crude ethanolic extract of *Peuraria Phaseoloide* Leaf (Roxb) Benth.(FABACEAE). *World Journal of Pharmaceutical Research*. Jun 1;10(7):39-59. Available at: <https://doi.org/10.20959/wjpr20217-20585>
  21. Olivia NU, Goodness UC, Obinna OM. (2021). Phytochemical profiling and GC-MS analysis of aqueous methanol fraction of *Hibiscus asper* leaves. *Future Journal of Pharmaceutical Sciences*. Dec;7:1-5. Available at: <https://doi.org/10.1186/s43094-021-00208-4>
  22. Hameed IH, Altameme HJ, Mohammed GJ. (2016). Evaluation of antifungal and antibacterial activity and analysis of bioactive phytochemical compounds of *Cinnamomum zeylanicum* (Cinnamon bark) using gas chromatography-mass spectrometry. *Oriental Journal of Chemistry*.32(4):1769. Available at: <http://dx.doi.org/10.13005/ojc/320406>
  23. Yu H, Feng J, Zhang Q, Xu H. (2015). (Z)-3-hexenyl acetate and 1-undecanol increase male attraction to sex pheromone trap in *Grapholita molesta* (Busck)(Lepidoptera: Tortricidae). *International Journal of Pest Management*. Jan 2;61(1):30-5. Available at: <https://doi.org/10.1080/09670874.2014.986560>
  24. Vanitha V, Vijayakumar S, Nilavukkarasi M, Punitha VN, Vidhya E, Praseetha PK. (2020). A novel microbiscidal bioactive alkane identified from *Plumbago zeylanica* L. *Industrial Crops and Products*. Oct 15;154:112748. Available at: <https://doi.org/10.1016/j.indcrop.2020.112748>
  25. Deep A, Phogat P, Kumar M, Kakkar S, Mittal SK, Malhotra M. (2012). New tetradecanoic acid hydrazones in the search for antifungal agents: synthesis and in vitro evaluations. *Acta Poloniae Pharmaceut Drug Res*. Jan 1;69:129-33.
  26. Umaru IJ, Badruddin FA, Umaru HA. Phytochemical, antifungal and antibacterial potential of *Leptadenia hastata* stem-bark extract. *MOJ Toxicol*. 2018;4(4):263-8.
  27. Aboh MI, Olayinka BO, Adeshina GO, Oladosu P. (2014). Antifungal activities of phyto compounds from *Mitracarpus villosus* (Sw.) DC from Abuja, Nigeria. *J Microbiol Res*.4(2):86-91. Available at: <https://doi.org/10.5923/j.microbiology.20140402.07>
  28. Umaru I, Fasihuddin A, Zaini B. (2017). Antifungal activity of *Leptadenia hastata* (Pers) Decne leaves extract. *Int. J. Pure Appl. Biosci*.5:14-8. Available at: <http://dx.doi.org/10.18782/2320-7051.5996>
  29. National Center for Biotechnology Information. (2023). PubChem Compound Summary for CID 31284, Methyl Tetradecanoate. Retrieved May 10. Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/Methyl-tetradecanoate>.
  30. Rao BG, Rao ES, Rao TM. (2012). Quantification of phytochemical constituents and in-vitro antioxidant activity of *Mesua ferrea* leaves. *Asian Pacific Journal of Tropical Biomedicine*. Feb 1;2(2):S539-42. Available at: [https://doi.org/10.1016/S2221-1691\(12\)60269-X](https://doi.org/10.1016/S2221-1691(12)60269-X)
  31. Imo C, Yakubu OE, Imo NG, Udegbonam IS, Onukwugha OJ. (2018). Chemical composition of *Xylopi aethiopica* fruits. *Am J Physiol Biochem Pharmacol*.7(2):48-53. Available at: <https://doi.org/10.5455/ajpbp.20180521064020>

32. Mahato SB, Sen S. (1997). Advances in triterpenoid research, 1990–1994. *Phytochemistry*. Apr 1;44(7):1185-236. Available at:[https://doi.org/10.1016/S0031-9422\(96\)00639-5](https://doi.org/10.1016/S0031-9422(96)00639-5)
33. Nestel P, Clifton P, Noakes M. (1994). Effects of increasing dietary palmitoleic acid compared with palmitic and oleic acids on plasma lipids of hypercholesterolemic men. *Journal of lipid research*. Apr 1;35(4):656-62. Available at: [https://doi.org/10.1016/S0022-2275\(20\)41179-4](https://doi.org/10.1016/S0022-2275(20)41179-4)
34. Shadrach P, Imo C, Otitoju O, Mohammad ZI, Asaph PM. (2023). Phytochemical Constituents of Ethanolic Extracts of Riped and Unripped Fruits of *Dennettia tripetala*. *Interna J of Pharma*02.5(1): 01-07.
35. Aparna V, Dileep KV, Mandal PK, Karthe P, Sadasivan C, Haridas M. (2012). Anti-inflammatory property of n-hexadecanoic acid: structural evidence and kinetic assessment. *Chemical biology & drug design*. Sep;80(3):434-9. Available at: <https://doi.org/10.1111/j.1747-0285.2012.01418.x>
36. Umaru I J, Ebenezer Morayo Ale, Kerenhappuch I U, Mulikat Adenike Adewole, Chinedu Christain Iheanacho, Glory Godwin Ebuk. (2024). Antibacterial Potential and Anti-Ulcer Activity of Methanol Extract of *Khaya anthotheca* on Indomethacin Induced Ulcer in Albino Wistar Rats. *Asian J. Biol. Sci.*, 17 (4): 496-504. <https://doi.org/10.3923/ajbs.2024.496.504>